MANTER: Journal of Parasite Biodiversity (ISSN 2470-8224)
Occasional Papers, Number 20, August 2, 2022
doi: 10.32873/unl.dc.manter20
https://digitalcommons.unl.edu/manter
Copyright © 2022 John E. Ubelaker and Gabor R. Racz

MANTER: Journal of Parasite Biodiversity

A Checklist of Parasites of *Peromyscus maniculatus* in North America

John E. Ubelaker[1] and Gabor R. Racz[2]

1 Emeritus Professor of Biological Sciences, Southern Methodist University, Dallas, Texas, USA
2 Manter Laboratory of Parasitology, University of Nebraska State Museum, University of Nebraska–Lincoln, Lincoln, Nebraska, USA 68588-0514

Abstract
This document represents a summary of parasites, in the broadest sense of the term, reported from *Peromyscus maniculatus* from throughout its range in North America. The document provides data from work ranging from paleontological findings to relatively current reports of parasites and parasitism from *P. maniculatus* and covers viruses, bacteria *sensu lato*, protists, helminths, and ectoparasites.

Keywords: parasites, *Peromyscus maniculatus*, virus, bacteria, ectoparasite, helminth parasite, protista, protozoa

Peromyscus maniculatus is a species of rodent, commonly known as the deer mouse, that belongs to the family Muridae Killiger, 1811. McKenna and Bell (1997) report that *P. maniculatus* belongs in the following classification:

Family Muridae Killiger, 1811
Subfamily Sigmodontinae Wagner, 1843
Tribe Peromyscini Hershkovitz 1966
Genus *Peromyscus* Gloger, 1841

Hibbard (1968) reports on the paleontology of *P. maniculatus* and cites the findings of Peterson (1926); Tamsitt (1957); Guilday (1962); Slaughter and Ritchie (1963); Guilday, Martin, and McCrady (1964); and Dalquest (1965). The oldest record is that of Slaughter and Ritchie with a Wisconsin ^{14}C age of 28,000 ± 4,740 BP years.

Peromyscus maniculatus is found throughout North America, from alpine regions to desert areas, from Alaska to central Mexico. The rodents are capable of living in every habitat in New Mexico but are most common in northern sage-grassland habitats. They are much less common in woodland, grassland, or desert communities. In southern New Mexico, the woodlands act as a barrier between lowland and mountain populations. In contrast, the mountain and lowland populations are in continuous contact throughout the woodland areas in northern New Mexico.

O'Farrell (1975) reported greater trapping success of deer mice at higher elevations, where they have a preference for more mesic *Artemisia*/*Agropyron* associations, with perennial grasses providing a source of succulent herbage as well as excellent cover and numerous additional habitats for insects. Richard (1960) found that deer mice became less abundant in the drier *Artemisia*/*Poa* habitats.

Populations of deer mice tend to be high but fluctuate throughout the year, with a peak in late summer or fall and a low in late winter. Gestation is from 21 to 28 days, and a litter varies from 1 to 10 with an average of 3 or 4 offspring. Weaning time is usually 25 to 35 days, and young mice become independent on average at 35 days. The average age of sexual maturity is 49 days. Holdenried and Morlan (1956) reported that pregnant mice in Santa Fe County, New Mexico, were encountered from December through September.

The mice are primarily nocturnal and spend their time on the ground within home ranges varying anywhere from 240 square meters to 3,000 square meters. They are om-

nivorous, consuming insects as well as seeds, fruit, flowers, and nuts (Jameson, 1952). Hamilton (1941) found some arthropods in the diet of deer mice in the eastern United States, and Whitaker Jr. (1966) also found arthropods in the diet of deer mice in Indiana. Koehler and Anderson (1991) found that this rodent relies heavily on arthropods for food during the months of April, June, and August in southern Idaho with lepidopteran larvae and carabid beetles occurring more frequently in their diets.

Numerous studies have provided evidence for territoriality or home ranges for *Peromyscus* (e.g., Burt, 1940; Nicholson, 1941; Redman and Sealander, 1958; Terman, 1961; Myton, 1974; Metzgar, 1971, 1973a, 1973b, 1979; Madison, 1977; Wolff, 1985a,b). According to Noble (1939), territories of *Peromyscus* are areas that are defended, whereas a home range is not. Territorial defense more likely occurs at high population densities rather than low. Sizes of home ranges or territories range from 790 to 3,000 square meters, and the variation is likely a function of habitat, food, and population density. Home ranges or territories are smaller in bushy, favorable habitats than in more open, less favorable habitats (Metzgar, 1971; Van Horne, 1982). Wolff (1985b) found that food does not appear to be the major factor that determines minimum size of home range, but males had slightly larger home ranges than females. Territoriality in *Peromyscus* appears to be the result of reproductive competition and not of defense of food (Nadeau et al., 1981; Wolff, 1985a,b). Males appear to be territorial to provide exclusive reproductive access to one or more females, whereas females probably defend a space to rear their young (Wolff and Lundy, 1985) or to prevent infanticide (Wolff, 1985b, 1993).

Articles about the parasites of *Peromyscus* species have been written by several authors. A review of the parasites of *Peromyscus* species has been written by Whitaker Jr. (1968), and King (1968) edited a special publication (no. 2) on the biology of *Peromyscus* (Rodentia). In this book, Whitaker Jr. wrote the chapter on parasites. Lichtenfels's (1966) thesis, written at the University of Maryland, reported on a survey of intestinal nematodes from Maryland rodents. Harkema (1936) wrote a monograph on the parasites of some North Carolina rodents. Hall (1916) wrote a monograph on the nematodes of Rodentia, Lagomorpha, and Hyracoidea. Doran (1954a,b, 1955a,b) presented the geographical distribution records of the protozoa and helminths of *P. maniculatus* in North America. Dyer (1969) provided a checklist of the protozoa and helminths of *P. maniculatus*. Mayberry et al. (2000) published a bibliography of parasites of vertebrate hosts in Arizona, New Mexico, and Texas that contains several references to parasites of *Peromyscus maniculatus*. Padovan (2006) summarized viral, bacterial, and fungal diseases associated with this species.

Because of its high population density and wide distribution, the deer mouse is considered responsible for distributing its parasites to other animals, especially other rodents, in all habitats and locations, as reported by Grundmann and Frandsen (1960).

Parasites

Viruses

Adenovirus (Adenoviridae) from Kern County, California, by Reeves et al. (1967), Hardy et al. (1974).

Buttonwillow virus (Bunyaviridae) from Kern County, California, by Taylor (1967, pp. 633–636) and Hardy et al. (1970).

Colorado tick fever virus (Reoviridae): experimental infection by Taylor (1967, pp 369–372) and Bowen et al. (1981a); natural infection reported by Taylor (1967), Burgdorfer (1960), Burgdorfer and Eklund (1960), Emmons (1988), Johnson (1970), Simpson (1968), Seymour and Yuill (1981), Bowen et al. (1981b), McLean et al. (1989), Clark et al. (1970), and Lane et al. (1982).

Hepatitis E virus (Hepesviridae) by Favorov et al. (2000).

Kobuvirus (Picornaviridae) by Phan et al. (2011).

Modoc virus (Flaviviridae) by Taylor (1967, pp 357–360) from Modoc County, California; Karabatsos (1985) and Hardy et al. (1974) from Kern County, California; Zarnke and Yuill (1985) from Alberta, Canada. Experimental infection by Davis et al. (1974) and Hume et al. (2015).

Monogahela virus (Bunyaviridae, Hantaviruses) by Song et al. (1996) and Monroe et al. (1999) in the eastern United States.

Mouse papillomavirus PmPV1 (Papillomaviridae) by Phan et al. (2011).

Powassan virus (Flaviviridae) by Taylor (1967, pp. 449–452) from the Black Hills, South Dakota; McDonald et al. (2009) from Alaska; Deardorff et al. (2013) from the southwestern US; Hardy et al. (1974) from California; Seymour and Yuill (1981).

Rodent adeno-associated virus (Parvoviridae) by Phan et al. (2011).

Rodent hepacivirus (Flaviviridae) by Kapoor et al. (2013).

Rodent stool-associated circular viruses RodSCV (Circoviridae) by Phan et al. (2011).

Saint Louis encephalitis virus (Flaviviridae) reported by Whitney et al. (1968) and by Seymour and Yuill (1981) based on serological diagnostic tests.

Sin Nombre virus (Bunyaviridae, Hantaviruses). Formerly known as Hantavirus Pulmonary Syndrome virus, Four Corners virus, Muerto Canyon virus, and Convict Creek virus. First reported by Nichol et al. (1993) from the

southwestern US. Additional report by Childs et al. (1994) in the southwestern United States; Elliot et al. (1994), Nerurkar et al. (1994), and Schmaljohn et al. (1994) in California; Graham and Chomel (1997) in California; Bennett et al. (1999) in California; Calisher et al. (1999) in Colorado; Feuer et al. (1999) in California and Nevada; Monroe et al. (1999) in the western US and Canada; Root et al. (1999) in Colorado; Boone et al. (2000) in California and Nevada; Drebot et al. (2001) in Canada; Glass et al. (2002) in the southwestern US; Kuenzi et al. (2005) in Montana; Douglas et al. (2007) in Montana; Madhav et al. (2007) in Montana; Safronetz et al. (2008) in Manitoba, Canada; Clay et al. (2009) in Utah; Disney et al. (2010) in Oregon; Carver et al. (2011) in Montana; Amman et al. (2013) in Colorado; and Danforth et al. (2020) in California.

Venezuelan equine encephalitis virus (Togaviridae) reported by Scherer et al. (1971), Sudia and Newhouse (1975), and Seymour and Yuill (1981) based on serological diagnostic tests.

Western equine encephalitis virus (Togaviridae) by Bowers et al. (1969) and Hardy et al. (1974).

Whitewater Arroyo virus (Arenaviridae). Woodrats (*Neotoma*) are regarded as the main reservoir of this virus. Bennett et al. (2000) used serological tests and found *P. maniculatus* blood to be reactive to this type of virus.

Bacteria

Anaplasma (*Erlichia*) *phagocytophilum* by Zeidner et al. (2000), DeNatale et al. (2002), and Larson et al. (2018).

Bartonella sp. by Tyzzer (1942), Kosoy et al. (1997), Jardine et al. (2005), Bai et al. (2011), and Goodrich et al. (2020).

Borrelia bissettii by Schneider et al. (2000), Burkot et al. (2000, 2001), DeNatale et al. (2002), Eisen et al. (2003, 2009), and Vredevoe et al. (2004).

Borrelia burgdorferi by Steere (1989), Rand et al. (1993), Ubico et al. (1996), Burkot et al. (1999), Zeidner et al. (2000), Oliver et al. (2006), Morshed et al. (2015), Larson et al. (2018), Buchholz et al. (2018), and Rodríguez-Rojas et al. (2020).

Coxiella burnetti by Stoenner et al. (1959), Marchette et al. (1962a), Vest et al. (1965), Enright et al. (1969, 1971a,b), and Thompson et al. (2012).

Francisella tularensis by Burroughs et al. (1945), Holdenried and Quan (1956), Vest and Marchette (1958), and Wobeser et al. (2007, 2009).

Leptospira sp. by Ferris et al. (1959), Redetzke and McCann (1980), and Verma et al. (2019).

Pasteurella multocida by Quan et al. (1979), but Swanekamp (2005) did not find it.

Rickettsia akari by Bennett et al. (2007).

Rickettsia rickettsii, causative agent of the Rocky Mountain Spotted Fever (RMSF), by Lundgren and Thorpe (1966) and Adjemian et al. (2008).

Yersenia pestis by Pollitzer and Meyer (1961), Kartman et al. (1962), Marchette et al. (1962b,c), Quan and Kartman (1962), Hudson et al. (1964, 1972), Hubbert et al. (1966), Hudson and Quan (1975), Nelson and Smith (1976), Twigg (1978), Barnes (1982), Poland et al. (1994), Davis et al. (2002), Nieto et al. (2007), Stapp et al. (2008), and Danforth et al. (2018). Experimental infection by Holdenried and Quan (1956) and Bacon and Drake (1958).

Fungi

Anthroderma benhamiae by Hubálek (2000).

Anthroderma persicolor by Hubálek (2000).

Anthroderma quadrifidum by Hubálek (2000).

Anthroderma silverae by Vanderwolf et al. (2018).

Coccidioides sp. by Catalán-Dibene et al. (2014), del Rocío Reyes-Montes (2016).

Emmonsia parva var.crescens is a saprotropic filamentous fungus associated with decaying plant material, but occasionally it can cause lung infection and disease. It appears that lung disease and pathology is caused by immune reaction to spores (Sigler, 2005), and this species is not a truly parasitic organism. Reported by Bakerspiegel (1956), Carmichael (1961), Dvorak et al. (1965), Jellison (1971). Spores reported in lung tissue by Bartlow (2019).

Microsporum versicolor by English et al. (1978), Hubálek and Rudolf (2011).

Trichophyton sp. by Carmichael (1961), Vanderwolf et al. (2018).

Protozoans

Besnoitia jellisoni from upper Salmon River, Idaho, by Frenkel and Lunde (1966).

Eimeria arizonensis from Oregon by Fuller (1996), from British Columbia by Levine and Ivens (1963), from New Mexico by Reduker et al. (1985).

Eimeria delicata from Oregon by Fuller (1996), from New Mexico by Reduker et al. (1985).

Eimeria peromysci from San Bernardino County, California, by Reduker et al. (1985); from New Mexico by Reduker et al. (1987).

Eimeria siniffi from British Columbia by Levine and Ivens (1963).

Giardia microti from California by Kofoid and Christiansen (1915a).

Giardia muris from California by Kofoid and Christiansen (1915b), Boeck and Stiles (1923).

Giardia peromysci from Ontario, Canada, by Grant and Woo (1979).
Grahamella sp. from Utah by Frandsen and Grundmann (1961b), from California by Wood (1975).
Haemobartonella sp. from California by Wood (1975).
Hepatozoon leptosoma from California by Wood (1962, 1975).
Isospora californica from White Sands National Monument, New Mexico, by Mayberry et al. (1980).
Isospora peromysci from northeastern Colorado and Ft. Collins, Colorado, by Duszynski and Anderson (1968); from California by Davis (1967).
Klossia perplexens from Illinois by Levine et al. (1955).
Sarcocystis idahoensis by Bledsoe (1979), Dubey (1983).
Sarcocystis peromysci by Dubey (1983).
Sarcocystis sp. from Benton County, Washington, by O'Farrell (1975).
Trichomonas muris from California by Kofoid and Christiansen (1915b), as cited by Doran (1955b).
Trypanosoma brucei experimental infection by Packchanian (1934).
Trypanosoma cruzi experimental infection by Wood (1934).
Trypanosoma peromysci from Alberta, Canada, by Watson and Hadwen (1912); from New Mexico, Arizona, and California by Wood (1942, 1952, 1975).
Trypanosoma sp. from Utah by Frandsen and Grundmann (1961b).

Arthropoda—Acari or mites acarines (Acarina)

Androlaelaps casalis by Allred (1957c), Allred and Beck (1966), Allred (1970), Whitaker and Maser (1985).
Androlaelaps debilis by Glicken and Schwab (1980).
Androlaelaps fahrenholzi (synonym of *Androlaeleps glasgowi*) by Allred (1957c, 1958), Allred and Beck (1966), Allred and Goates (1964), Douglas (1969), Elzinga and Rees (1964), Glicken and Schwab (1980), Hansen (1964), Keegan (1953), Lawrence et al. (1965), Rapp (1962), Reeves et al. (2007), Scholten et al. (1962), Storm and Ritzi (2008), Whitaker and Corthum (1967), Whitaker et al. (1975), Whitaker and Maser (1985), Wyman and Schaefer (1972), Howell et al. (2016).
Androlaelaps leviculus by Allred (1957c), Allred (1958). Allred and Beck (1966), Keegan (1951), Keegan (1956).
Anoetidae sp. by Whitaker and Maser (1985).
Antennophorid sp. by Allred (1957c).
Ascaidae sp. by Allred (1957c).
Bdellonyssus bacoti from Ft. Hood, Texas, by Hedeen (1953).
Brevisterna morlani by Strandtmann and Allred (1956).
Brevisterna utahensis by Stradtmann and Allred (1956), Allred (1957c), Allred and Beck (1966).
Camisiidae sp. by Allred (1957c).
Cheyletidae sp. by Allred (1957c).
Cyrtolaelaps sp. by Whitaker and Maser (1985).
Dermacarus hypudaei by McDaniel (1979), Rupes and Whitaker Jr. (1968), Whitaker et al. (1975).
Dermacarus jonesi by Fain and Whitaker Jr. (1976).
Dermacarus newyorkensis by Fain et al. (1971).
Dermanyssus becki by Allred (1957c), Allred and Beck (1966).
Eremaeidae sp. by Allred (1957c).
Eubrachylaelaps circularis by Allred (1957c), Allred and Beck (1966), Furman (1955), Jameson Jr. and Brennan (1957), Whitaker and Maser (1985).
Eubrachylaelaps debilis by Allred (1957c), Allred and Beck (1966), Jameson Jr. and Brennan (1957), Keegan (1953), Whitaker and Maser (1985).
Eubrachylaeleps hollisteri by Allred (1957c), Allred and Beck (1966).
Eulaelaps sp. by Allred (1957c).
Eulaelaps stabularis by Allred and Beck (1966), Hansen (1964), Lawrence et al. (1965), Whitaker Jr. and Wilson (1968), Whitaker et al. (1975), Whitaker and Maser (1985).
Euryparasitus sp. by Whitaker and Maser (1985).
Gamasolaelaptidae sp. by Allred (1957c).
Garmania sp. by Allred (1957c).
Glycyphagus hypudaei by Whitaker and Maser (1985), Reeves et al. (2007), Storm and Ritzi (2008).
Glycyphagus sp. by Allred (1957c).
Haemogamasus ambulans by Allred (1957c), Allred and Beck (1966), Keegan (1951), Wyman and Schaefer (1972).
Haemogamasus liponyssoides by Keegan (1951), Lawrence et al. (1965), Storm and Ritzi (2008).
Haemogamasus longitarsus by Allred (1957c), Allred and Beck (1966).
Haemogamasus nidi by Allred (1957c), Allred and Beck (1966), Hansen (1964).
Haemogamasus pontiger by Allred (1957c), Allred and Beck (1966).
Haemogamasus reidi by Whitaker and Maser (1985).
Haemogamasus sp. by Allred (1957c).
Hirstionyssus blarinae by Storm and Ritzi (2008), cited as *Echinonyssus blarinae*.
Hirstionyssus carnifex by Keegan (1953).
Hirstionynssus cynomys by Rapp (1962).
Hirstionynssus geomydis by Allred (1957c).
Hirstionyssus hilli by Allred (1957c), Allred and Beck (1966), Herrin (1970), Keegan (1953).
Hirstionyssus incomptis by Allred (1957c), Allred and Beck (1966), Elzinga and Rees (1964), Herrin (1970).
Hirstionyssus isabellinus by Allred (1957c), Allred and Beck (1966).
Hirstionyssus neotomae by Allred and Beck (1966).

Hirstionyssus obsoletus by Allred (1957c), Jameson (1950), and by Whitaker and Maser (1985), cited as *Echinonyssus obsoletus*.
Hirstionyssus occidentalis by Elzinga and Rees (1964), Hansen (1964), Herrin (1970), Strandtmann and Morlan (1953).
Hirstonyssus punctatus by Allred and Beck (1966).
Hirstionyssus sp. by, and perhaps species "p," Scholten et al. (1962) and Johnson and Brennan (1957).
Hirstionyssus talpae by Hansen (1964), Whitaker Jr. and Wilson (1968).
Hirstionyssus tarsalis by Allred and Beck (1966).
Hirstionyssus triacanthus by Allred and Beck (1966).
Hirstionyssus utahensis by Allred and Beck (1966), Glicken and Schwab (1980), Herrin (1970), Whitaker et al. (1975). By Storm and Ritzi (2008), Whitaker and Maser (1985), cited as *Echinonyssus utahensis*.
Hypoaspis lubrica by Allred (1957c), Allred and Beck (1966), Storm and Ritzi (2008).
Hypoaspis miles by Allred (1957c), Allred and Beck (1966).
Hypoaspis sp. by Allred (1957c), Whitaker et al. (1975).
Ischyropoda armatus by Allred (1957c), Allred and Beck (1966), Keegan (1951).
Ischylopoda furmani by Allred (1965), Allred and Beck (1966), Keegan (1956).
Kleemania sp. by Allred (1957c), Allred and Goats (1964), Whitaker Jr. and Wilson (1968).
Labidophoridae sp. by Scholten et al. (1962).
Laelaps incilis by Allred and Beck (1966).
Laelaps kochi by Whitaker Jr. and Wilson (1968), Whitaker and Maser (1985).
Laelaps nuttalli by Allred (1957c), Allred and Beck (1966), Ulrich and Vaughn (1963).
Listrophorus sp. by Allred (1957c), Allred and Beck (1966).
Macrocheles sp. by Allred and Beck (1966), Hansen (1964).
Macrochelidae sp. By Whitaker Jr. and Wilson (1968).
Myobia musculi by McDaniel (1979), Storm and Ritzi (2008), Ulrich and Vaughn (1963).
Myobiidae sp. by Scholten et al. (1962).
Myocoptes japonensis by Storm and Ritzi (2008), Whitaker et al. (1975).
Myocoptes musculinus by McDaniel (1979), Reeves et al. (2007), Storm and Ritzi (2008).
Myocoptes sp. by Allred and Beck (1966), Elzinga and Rees (1964).
Neoparasitidae sp. by Allred (1957).
Ondatralaelaps multispinosus by Allred (1957c), Allred and Beck (1966).
Ornithonyssus bacoti by Allred (1957c), Allred and Beck (1966), Connior et al. (2017), Whitaker Jr. and Wilson (1968).
Ornithonyssus sp. by Allred (1957c).
Orycteroxenus soricis by Storm and Ritzi (2008).
Pachylaelapidae sp. by Allred (1957c).
Parasitidae sp. by Allred (1957c).
Phytoseiidae sp. by Allred (1957c).
Poecilochiridae sp. by Whitaker Jr. and Wilson (1968).
Poecilochirus sp. by Allred (1957c), Hansen (1964).
Proctolaelaps sp. by Whitaker and Maser (1985).
Protomyobia claparedei by Whitaker et al. (1975).
Psorergates peromysci by Giesen et al. (1983).
Psorergates watsoni by Kok et al, (1971).
Pyemotidae sp. by Allred (1957c).
Pygmephorus sp. by Hansen (1964).
Radfordia affinis by Whitaker Jr. and Wilson (1968).
Radfordia bachai by Howell and Elzinga (1962).
Radfordia lemnina by Allred (1957c), Allred and Beck (1966).
Radfordia subuliger by Allred (1957c), Allred and Beck (1966), Elzinga and Rees (1964), Fain and Bochkov (2002), Reeves et al. (2007), Spencer (1940), Storm and Ritzi (2008), Whitaker Jr. and Wilson (1968), Whitaker et al. (1975).
Resinacarus sp. by Whitaker Jr. and Wilson (1968).
Rhizoglyphus echinopus by Allred (1957c).
Tetranychidae sp. by Allred (1957c).
Trichoecius tenax by Storm and Ritzi (2008).
Typhlodromus mariposus by Allred (1957c).
Uropodidae sp. by Allred (1957c).

Acarina—Trombiculidae

Chatia ochotona by Allred (1957c), Allred and Beck (1966), Brennan and Beck (1955), Gould (1956), Traub and Nadchatram (1966).
Chatia setosa by Allred (1957c), Brennan (1946), Brennan and Beck (1955), Easton (1975), Gould (1956), Jameson Jr. and Brennan (1957), Traub and Nadchatram (1966).
Cheladonta micheneri by Loomis (1956).
Comatacarus inconspicuus by Reed (1973).
Comatacarus pusillus by Goff and Loomis (1974).
Euschoengastia ambocalis by Wrenn and Loomis (1973).
Euschoengastia californica by Gould (1956), Loomis and Bunnell (1962).
Euschoengastia cordiremus by Allred and Beck (1966), Allred and Goates (1964), Brennan (1948), Brennan and Beck (1955), Gould (1956).
Euschoengastia crateris by Farrell (1956).
Euschoengastia criceticola by Allred (1957c), Allred and Beck (1966), Allred and Goates (1964), Brennan (1948), Brennan and Beck (1955), Brennan and Jones (1954), Brown and Brennan (1952), Douglas (1969), Elzinga and Rees (1964), Gould (1956), Jameson and Brennan (1957).
Euschoengastia decipiens by Allred and Beck (1966), Brennan and Beck (1955).
Euschoengastia diverse by Loomis (1956).
Euschoengastia frondifera by Gould (1956).

Euschoengastia jamesoni by Whitaker et al. (1975)
Euschoengastia jonesi by Lipovsky and Loomis (1954), Loomis (1956)
Euschoengastia lacerta by Gould (1956)
Euschoengastia lanei by Allred and Beck (1966), Allred and Goates (1964), Douglas (1969).
Euschoengastia luteodema by Allred and Beck (1966), Brennan and Beck (1955).
Euschoengastia obesa by Allred and Beck (1966), Brennan and Beck (1955).
Euschoengastia oregonensis by Allred and Beck (1966), Brennan and Beck (1955), Easton (1975).
Euschoengastia peromysci by Brennan and Jones (1954), Douglas (1969), Farrell (1956), Gould (1956), Jameson Jr. and Brennan (1957), Loomis (1956), Reeves et al. (2007), Storm and Ritzi (2008), Whitaker et al. (1975).
Euschoengastia pomerantzi by Allred and Beck (1966), Gould (1956).
Euschoengastia radfordi by Allred and Beck (1966), Brennan and Jones (1954), Elzinga and Rees (1964), Gould (1956), Jameson Jr. and Brennan (1957).
Euschoengastia rotunda by Allred and Beck (1966).
Euschoengastia sciuricola by Allred (1957c), Allred and Beck (1966), Gould (1956), Jameson Jr. and Brennan (1957).
Euschoengastia setosa by Lawrence et al. (1965), Loomis (1956), Reeves et al. (2007), Whitaker et al. (1975).
Euschoengastia trigenuala by Loomis (1956).
Euschoengastia velata by Easton (1975).
Euschoengastoides hoplai by Brennan and Jones (1954), Gould (1956).
Euschoengastoides loomisi by Crossley and Lipovsky (1954), Loomis (1956).
Euschoengastoides neotomae by Loomis (1971).
Eutrombicula alfreddugesi by Loomis (1956), Wolfenbarger (1952).
Eutrombicula lipovskyana by Loomis (1956).
Fonsecia gurneyi by Loomis (1955, 1956).
Fonsecia kansasensis by Loomis (1955, 1956).
Hyponeocula arenicola by Tanigoshi and Loomis (1974).
Leeuwenhoekia americana by Allred and Beck (1966), Gould (1956), Loomis (1956). Reed (1973) revised the genus and cites this species as *Comatacarus americanus*.
Leeuwenhoekia dolosa by Gould (1956), Jameson and Brennan (1957). Reed (1973) revised the genus and cites this species as *Comatacarus dolosus*.
Leewenhoekia (*Comatacarus*) *americanus* by Douglas (1969), Easton (1975).
Leptotrombidium myotis by Allred (1957c), Allred and Beck (1966), Lawrence et al. (1965).
Leptotrombidium panamensis by Allred and Beck (1966).
Microtrombicula crossleyi by Loomis (1954), Vercammen-Grandjean (1965).
Microtrombicula nasalis by Webb and Loomis (1970).
Microtrombicula ornata by Loomis (1956), Loomis and Lipovsky (1954), Webb and Loomis (1970).
Miyatrombicula esoensis by Allred (1957c), Allred and Beck (1966).
Miyatrombicula scottae by Brennan and Jones (1954), Gould (1956).
Miyatrombicula sp. by Lawrence et al. (1965).
Neotrombicula brennani by Easton (1975).
Neotrombicula browni by Brennan and Wharton (1950), Kardos (1954).
Neotrombicula californica by Allred (1957c), Allred and Beck (1956), Brennan and Beck (1955), Brennan and Wharton (1950), Gould (1956).
Neotrombicula cavicola by Brennan and Wharton (1950), Gould (1956).
Neotrombicula carterae by Withaker et al. (1975).
Neotrombicula dinehartae by Brennan and Jones (1954), Brennan and Wharton (1950), Gould (1956).
Neotrombicula harperi by Allred (1957c), Allred and Beck (1966), Brennan and Wharton (1950), Kardos (1954), Lawrence et al. (1965).
Neotrombicula jewetti by Allred (1957c), Allred and Beck (1966), Brennan and Jones (1954), Brennan and Wharton (1950), Gould (1956), Jameson Jr. and Brennan (1957).
Neotrombicula lipovskyi by Kardos (1954), Loomis (1956).
Neotrombicula loomisi by Kardos (1954), by Douglas (1969) as *Trombicula loomisi*.
Neotrombicula microti by Allred (1957c), Brennan and Wharton (1950), Brown and Brennan (1952), Jameson Jr. and Brennan (1957).
Neotrombicula sylvilagi by Kardos (1954), Loomis (1956).
Neotrombicula whartoni by Kardos (1954).
Odontacarus dentatus by Goff et al. (1972), used as a senior synonym of *O. galli*.
Odontacarus galli by Greenberg (1952), Loomis (1956).
Odontacarus hirsutus by Jameson Jr. and Brennan (1957).
Odontacarus linsdalei by Allred and Beck (1966), Allred and Goates (1964).
Odontacarus villosus by Goff and Loomis (1973).
Pseudoschoengastia farneri by Lipovsky (1951a), Loomis (1956).
Pseudoschoengastia hungerfordi by Lipovsky (1951a, 1951b), Loomis (1956).
Pseudoschoengastia occidentalis by Loomis and Bunnell (1962).
Trombicula arenicolor by Allred and Beck (1966), Brennan and Beck (1955), Elzinga and Rees (1964).
Trombicula bakeri by Allred and Beck (1966).
Trombicula montanensis by Allred (1957c), Allred and Beck (1966).
Xenodontacarus plumosus by Loomis and Goff (1973); by Greenberg (1951) and Loomis (1956), cited as *Odontacarus plumosus*.

Anoplura

Hoplopleura acanthopus by Augustson (1941), Hansen (1964), Lawrence et al. (1965).

Hoplopleura erratica by Lawrence et al. (1965).

Hoplopleura hesperomydis from California by Ferris (1921), Augustson (1941), Cook (1959), Elzinga and Rees (1964), Ferris (1916), Glicken and Schwab (1980), Hansen (1961), Kellogg and Ferris (1915), Kim (1965), Kim et al. (1966), Lawrence et al. (1965), Morlan and Hoff (1957), Reeves et al. (2007), Scanlon (1960), Scholten et al. (1962), Spencer (1966), Storm and Ritzi (2008), Whitaker et al. (1975), Wilson (1961), Wyman and Schaefer (1972), Zwolak et al. (2013).

Hoplopleura sciuiricola by Lawrence et al. (1965).

Neohaematopinus laeviusculus by Augustson (1941).

Polyplax auricularis by Augustson (1941), Beer et al. (1959), Morlan and Hoff (1957), Elzinga and Rees (1964), Emerson (1971), Ferris (1916, 1951), Hansen (1964), Kellogg and Ferris (1915, 1916), Reeves et al. (2007), Scanlon (1960), Spencer (1966), Whitaker et al. (1975), Zwolak et al. (2013).

Polyplax serrata by Ulrich and Vaughn (1963).

Ticks

Amblyomma maculatum by Semtner and Hair (1973).

Dermacentor andersoni from Colorado, 0.59 larvae, 4.47 nymphs by Carey et al. (1980); Clark et al. (1970), Grundmann and Frandsen (1960), Augustson (1941a), Douglas (1969), Hansen (1964), Johnson (1966), Hunter and Bishopp (1941), Sonenshine et al. (1976), Zwolak et al. (2013).

Dermacentor occidentalis by Coultrip et al. (1973), Bishopp and Trembley (1945), Holdenried et al. (1951), Lane et al. (1981), Mohr et al. (1964), Wyman and Schaefer (1972).

Dermacentor parumapertus by Beck et al. (1963), Gastfriend (1955), Johnson (1966).

Dermacentor sp. by Bacon et al. (1959), Coultrip et al. (1973).

Dermacentor variabilis by Campbell and MacKay (1979), Dodds et al. (1969), Garvie et al (1978), Lawrence et al. (1965), Reeves et al. (2007), Whitaker et al. (1975).

Ixodes angustus by Allred et al. (1960), Beck, et al. (1963), Bishopp and Trembley (1945), Cooley and Kohls (1945), Douglas (1969), Easton and Goulding (1974), Fay and Rausch (1969), Johnson (1966), Lawrence et al. (1965), Reeves et al. (2007), Whitaker et al. (1975), Zwolak et al. (2013).

Ixodes cookei by Cooley and Kohls (1945).

Ixodes kingi by Allred et al. (1960), Beck et al. (1963), Johnson (1966), Sidwell et al. (1964).

Ixodes marmoratae by Allred et al. (1960).

Ixodes muris by Allred et al. (1960).

Ixodes ochotonae by Allred et al. (1960), Clark et al. (1970), Johnson (1966).

Ixodes pacificus by Allred et al. (1960), Easton and Goulding (1974), Holdenried et al. (1951), Johnson (1966), Sidwell et al. (1964).

Ixodes peromysci by Cooley and Kohls (1945).

Ixodes sculptus by Bishopp and Trembley (1945).

Ixodes sp. by Allred et al. (1960), Coultrip et al. (1973), Hansen (1964), Holdenried et al. (1951), Scholten et al. (1962).

Ixodes spinipalpis by Allred et al. (1960), Beck (1955), Douglas (1969), Johnson (1966), Mohr et al. (1964).

Ixodes woodi from Texas by Eads et al. (1956).

Ornithodoros eremicus by Cooley and Kohls (1941).

Otobius sparnus by Johnson (1966), Kohls and Clifford (1963); Kohls et al. (1965) cited as *Ornithodoros sparnus*.

Diptera

Cuterebra americana complex by Grundmann and Frandsen (1960).

Cuterebra approximata by Smith (1975, 1977).

Cuterebra cyanella by Douglas (1969).

Cuterebra grisea from British Columbia by Hunter et al. (1972).

Cuterebra larvae from the Great Basin by Llewellyn (1978).

Cuterebra sp. by Blair (1942), Lawrence et al. (1965), Maurer and Skaley (1968), Scholten et al. (1962), Seaman and Nash (1976).

Fleas

Aetheca thamba (Jordan, 1929) by Holland (1944), Kinsella and Pattie (1967), Kinsella (1968), all cited as *Monopsyllus thambus*.

Aetheca wagneri Baker, 1904 from Bernalillo, Rio Arriba, and Sandoval County, New Mexico, by Ford et al. (2004); from Montana by Zwolak et al. (2013); from Utah by Stanford (1931) cited as *Ceratophyllus wagneri*.

Amalareus penicilliger (Grube, 1851) by Hubbard (1947), Jordan (1932), both cited as *Malaraeus penicilliger*.

Amaradix bitterrootensis Dunn and Parker, 1923 by Augustson (1941a), Hubbard (1947), Stark (1958), all cited as *Malaraeus bitterrootensis*.

Amaradix euphorbi (Rothschild, 1905) by Baker (1905), Beck and Allred (1966), Allred (1968), Egoscue (1966, 1976), Elzinga and Rees (1964), Haas et al. (1973), Hansen (1964), Holland (1949), Hubbard (1947), Morlan (1955), Senger (1966), Stark (1958), Tipton and Mendez (1968), Wagner (1936), Woods and Larson (1971), all cited as *Malaraeus euphorbi*; by Ford et al. (2004), Zwolak et al. (2013).

Amphipsylla sibirica (Wagner, 1898) by Allred (1968), Eads et al. (1979).

Anomiopsyllus amphibolus Wagner, 1936 by Barnes et al. (1977), Beck and Allred (1966), Egoscue (1976), Stark (1958).

Anomiopsyllus falsicalifornicus C. Fox 1929 by Coultrip et al. (1973).

Anomiopsyllus hiemalis mexicanus Holland, 1965 from Santa Fe County, New Mexico, by Ford et al. (2004).

Anomiopsyllus novomexicanensis Williams and Hoff, 1951 by Barnes et al. (1977); from Bernalillo and Hidalgo Counties, New Mexico, by Ford et al. (2004).

Anomiopsyllus nudatus (Baker, 1898) by Barnes et al. (1977), Haas et al. (1973), Stark (1958).

Atyphloceras echis Jordan and Rothschild, 1915 from Hidalgo, New Mexico, by Ford et al. (2004); by Egoscue (1976), Stark (1958).

Atyphloceras multidentatus (C. Fox, 1909) from Oregon by Easton (1983), Jameson and Brennan (1957), Rutledge et al. (1979), Stark (1958), Stark and Kinney (1969); by Holland (1949) as *A. artius* Jordan, 1933; by Hubbard (1947) as *A. felix* Jordan, 1933.

Callistopsyllus terinus Rothschild, 1905 from Santa Fe County, New Mexico, by Ford et al. (2004), Allred (1968), Egoscue (1976), Hansen (1964), Holland (1941, 1949), Hubbard (1947, 1949), Morlan (1955), Senger (1966); by Augustson (1941a), Beck (1966), Douglas (1969), and Stark and Kinney (1969) as *C. deuterus* Jordan, 1937.

Carteretta carteri Fox, 1927 by Holdenried et al. (1951), Hubbard (1947).

Catallagia calisheri Eads and Campos, 1979 by Eads and Campos (1979).

Catallagia charlottensis (Baker, 1898) by Baker (1905), Dunn and Parker (1923), Wagner (1936), and Hubbard (1947), who cites it both as *C. charlottensis* and as *C. motei* Hubbard, 1940.

Catallagia decipiens Rothschild, 1915 from Rio Arriba, Sandoval, Santa Fe County, New Mexico, by Ford et al (2004); from Benton County, Washington, by O'Farrell (1975), Allred (1952), Allred (1968), Beck (1966), Douglas (1969), Egoscue (1966, 1976), Haas et al. (1973), Hansen (1964), Holland (1949), Hubbard (1947, 1949), Kinsella (1968), Morlan (1955), Zwolak et al. (2013).

Catallagia mathesoni Jameson, 1950 by Jameson and Brennan (1957), Nelson and Smith (1976), Stark and Kinney (1969).

Catallagia sculleni Hubbard 1940 by Hubbard (1947), Jameson and Brennan (1957), Nelson and Smith (1976), Poinar and Nelson (1973), Stark and Kinney (1969); from Oregon by Easton (1983), Holdenried et al. (1951) as *C. chamberlini* Hubbard, 1940; from California Augustson (1941b), both as *C. rutherfordi* and *C. vonbloekeri*.

Cediopsylla inaequalis (Baker, 1895) by Allred (1968).

Ceratophyllus ciliatus Baker, 1904 from Oregon by Easton (1983); by Holland (1941), Hubbard (1947), Jameson et al. (1957), all cited as *Monopsyllus ciliatus*.

Ceratophyllus gallinae (Schrank, 1803) by Reeves et al. (2007).

Ceratophyllus pelecani Augustson 1942 by Hubbard (1947).

Conorhinopsylla nidicola Jellison, 1945 by Poorbraugh and Gier (1961).

Corrodopsylla curvata Rothschild, 1915 by Hansen (1964); by Hubbard (1947) as *Doratopsylla jellisoni* Hubbard, 1940; by Robert (1962).

Corypsylla jordani Hubbard, 1940 by Hubbard (1947).

Corypsylla ornata C. Fox, 1908 by Hubbard (1947).

Ctenophthalmus pseudagyrtes Baker, 1904 by Benton and Krug (1956), Benton et al. (1969), Buckner (1964), Geary (1959), Holland (1949), Lawrence et al. (1965), Layne (1958), Main (1970), Poorbraugh and Gier (1961), Robert (1962), Scharf and Stewart (1980), Verts (1961), Whitaker and Corthum (1967), Whitaker et al. (1975), Woods and Larson (1971).

Dasypsyllus gallinulae (Dale, 1878), invasive species by Easton (1983).

Delotelis hollandi Smit, 1952 by Jameson and Brennan (1957).

Delotelis telegoni (Rothschild, 1905) by Augustson (1941a), Holland (1949), Hubbard (1947).

Doratopsylla blarinae C. Fox, 1914 from northern New York state by Benton et al. (1969).

Doratopsylla jellison, synonym of *Corrodopsylla curvata*.

Echidnophaga gallinacea (Westwood, 1875) by Beer et al. (1959), Hardy et al. (1974).

Epitedia scapani (Wagner, 1936) from California by Jameson and Brennan (1957); from Oregon by Easton (1983), Holland (1941, 1949), Hubbard (1947), cited as *E. jordani* Hubbard, 1940.

Epitedia stanfordi Traub, 1944 from Sandoval and Santa Fe Counties, New Mexicio, by Ford et al. (2004); by Allred (1968), Beck (1966), Douglas (1969), Egoscue (1976), Elzinga and Rees (1964), Hubbard (1947), Morlan (1955), Stark (1958).

Epitedia wenmanni Rothschild, 1904 from Rio Arriba by Ford et al. (2004), Allred (1952), Allred (1968), Augustson (1941a), Beer et al. (1959), Benton and Krug (1956), Benton and Altmann (1964), Benton et al. (1969), Buckner (1964), Dunn and Parker (1923), Fox (1940), Geary (1959), Holland (1949), Hubbard (1947), Jameson and Brennan (1957), Lawrence et al. (1965), Layne (1958), Main (1970), Poorbaugh and Gier (1961), Rapp and Gates (1957), Reeves et al. (2007), Robert (1962), Scharf and Stewart (1980), Scholten et al. (1962), Schwan and Dobkin (1981), Stark (1958), Storm and Ritzi (2008), Tipton and Mendez (1968), Verts (1961), Ulrich and Vaughn (1963), Whitaker and Corthum (1967), Whitaker et al. (1975), Wilson (1957), Wyman and Schaefer (1972), Zwolak et al. (2013).

Eumolpianus eumolpi Rothschild, 1905 by Allred (1952), Allred (1968), Beck and Allred (1966), Haas et al. (1973), Hansen (1964), Hubbard (1947), Morlan (1955), Sidwell

et al. (1964), Stark (1958), Stark and Kinney (1969), all citing it as *Monopsyllus eumolpi*. Also by Ford et al. (2004) from Sandoval County, New Mexico.

Foxella ignota (Baker, 1895) from Sandoval County, New Mexico, by Ford et al (2004), Allred (1952), Allred (1968), Buckner and Blasko (1969), Hubbard (1947), Stark (1958). By Haas et al. (1973) cited as *Dactylopsylla ignota*.

Hoplopsyllus anomalus (Baker, 1904) by Hubbard (1941).

Hystrichopsylla dippiei Rothschild, 1902 from Sandoval County, New Mexico, by Ford et al. (2004), Beer et al. (1959), Buckner (1964), Egoscue (1976), Haas et al. (1973), Holland (1949), Scharf and Stewart (1980), Senger (1966), Stark (1958), while Allred (1952) and Hubbard (1947, 1949) cited as *H. gigas dippiei* (Baker, 1895).

Hystrichopsylla occidentalis Holland, 1949 from Oregon by Easton (1983); from Alaska, California, and Colorado by Campos and Stark (1979), Hubbard (1947); from Utah by Allred (1968). By Egoscue (1966, 1976) cited as *Hystrichopsylla linsdalei* Holland 1957.

Leptopsylla segnis (Schonherr, 1811) by Hardy et al. (1974).

Malaraeus sinomus (Jordan, 1925) from Bernalillo, Sandoval, and Santa Fe Counties, New Mexico, by Ford et al (2004); from Utah by Stanford (1931), cited as *Ceratophyllus sinomus*; by Beck and Allred (1966), Beck (1966), Douglas (1969), Egoscue (1976), Haas et al. (1973), Hardy et al. (1974), Hubbard (1947, 1949), Morlan (1955), Stanford (1931), Sidwell et al. (1964), Stark (1958).

Malareus telchinus (Rothschild, 1905) from Rio Arriba, Sandoval, and Santa Fe Counties, New Mexico, by Ford et al. (2004); from Montana by Dunn and Parker (1923); from Fresno County, California, by Hubbard (1941); from Oregon by Easton (1983). By Allred (1952), Allred (1968), Beck and Allred (1966), Beck (1966), Coultrip et al. (1973), Douglas (1969), Dunn and Parker (1923), Egoscue (1976), Glicken and Schwab (1980), Haas et al. (1973), Hansen (1964), Hardy et al. (1974), Holdenried et al. (1951), Holland (1949), Hubbard (1947, 1949), Hubbard (1941), Jameson Jr. and Brennan (1957), Morlan (1955), Rutledge et al. (1979), Stark (1958), Stark and Kinney (1969), Wheldon (1941), Zwolak et al. (2013).

Megabothris abantis (Rothschild, 1905) from Rio Arriba and Sandoval Counties by Ford et al (2004); from Oregon by Easton (1983). By Allred (1952); Augustson (1941a); Egoscue (1976); Haas et al. (1973); Hansen (1964); Holland (1949); Hubbard (1947), who also cites it as *M. adversus* (Wagner, 1936); Kinsella and Pattie (1967); Stark (1958); Zwolak et al. (2013).

Megabothris asio (Baker, 1904) by Holland (1949), Lawrence et al. (1965), Wilson (1961).

Megabothris lucifer (Rothschild, 1905) by Holland (1949).

Megabothris quirini (Rothschild, 1905) by Benton et al. (1969), Benton et al. (1971), Buckner (1964), Holland (1949), Lawrence et al. (1965), Robert (1962), Scharf and Stewart (1980).

Megarthroglossus bisetis Jordan and Rothschild, 1915 from Bernalillo and Sandoval Counties, New Mexico, by Ford et al. (2004), Williams and Hoff (1951).

Megarthroglossus divisus (Baker, 1898) from Bernalillo and Santa Fe Counties, New Mexico, by Ford et al. (2004). By Holland (1949), Kinsella and Pattie (1967), Morlan (1955).

Megarthroglossus procus Jordan and Rothschild, 1915 by Augustson (1941a), Beck (1966), Douglas (1969), Egoscue (1976), Hubbard (1947), Jameson and Brennan (1957).

Megarthroglossus smiti Medez, 1956 by Egoscue (1976).

Megarthroglossus weaveri Eads and Campos, 1977 by Eads and Campos (1977).

Meringis altipecten Traub and Hoff, 1951 from Hidalgo County, New Mexico, by Ford et al. (2004).

Meringis arachis Jordan, 1929 from Hidalgo County, New Mexico, by Ford et al. (2004).

Meringis bilsingi Eads and Menzies, 1949 from Doña Ana, New Mexico, by Ford et al. (2004).

Meringis cummingi (C. Fox, 1926) by Coultrip et al. (1973), Hubbard (1947), Hardy et al. (1974).

Meringis dipodomys Kohls, 1938 by Egoscue (1976), Hubbard (1947), Stark (1958).

Meringis hubbardi Kohls, 1938 by Allred (1968), Egoscue (1966, 1976), Hansen (1964), Hubbard (1947).

Meringis nidi Williams and Hoff, 1951 from Santa Fe County, New Mexico, by Ford et al. (2004). By Morlan (1955).

Meringis parkeri Jordan, 1937 from Sandoval and Santa Fe Counties, New Mexico, by Ford et al. (2004). By Allred (1968), Beck and Allred (1966), Haas et al. (1973), Hansen (1964), Hardy et al. (1974), Hubbard (1947), Morlan (1955), Sidwell et al. (1964), Stark (1958), Stark and Kinney (1969), Wyman and Schaefer (1972).

Meringis rectus Morlan, 1953 from Santa Fe County, New Mexico, by Ford et al. (2004). By Morlan (1955).

Meringis shannoni (Jordan, 1929) from Benton County, Washington, by O'Farrell (1975). By Holland (1949), Hubbard (1947).

Monopsyllus vison (Baker, 1904) by Benton et al. (1969), Holland (1949).

Monopsyllus wagneri (Baker, 1904) from Benton County, Washington, by O'Farrell (1975); from Oregon by Easton (1983); from California by Poiner and Nelson (1973). By Allred (1952, 1968), Augustson (1941a), Beck and Allred (1966), Beck (1966), Beer et al. (1959), Brown (1944), Buckner (1964), Coultrip et al. (1973), Douglas (1969), Dunn and Parker (1923), Egoscue (1976), Elzinga and Rees (1964), Glicken and Schwab (1980), Haas et al. (1973), Hansen (1964), Hardy et al. (1974), Holdenried

et al. (1951), Holdenried and Morlan (1955, 1956), Holland (1949), Hubbard (1947, 1949), Jameson and Brennan (1957), Jordan (1929), Kinsella and Pattie (1967), Kinsella (1968), Lawrence et al. (1965), Morlan (1955), Nelson and Smith (1976), Poinar and Nelson (1973), Rapp and Gates (1957), Rutledge et al. (1979), Schwan and Dobkin (1981), Sidwell et al. (1964), Stanford (1931), Stark (1958), Stark and Kinney (1969), Stark et al. (1976), Ulrich and Vaughn (1963), Verts (1961), Wagner (1936), Woods and Larson (1970), Wyman and Schaefer (1972).

Nearctopsylla hamata Holland and Jameson, 1949 by Jameson and Brennan (1957).

Nearctopsylla princei Holland and Jameson, 1949 by Jameson and Brennan (1957).

Neopsylla inopina Rothschild, 1915 by Hubbard (1949).

Nosopsyllus fasciatus (Bosc, 1800) by Allred (1952), Emmons et al. (1970), Holland (1949), Hubbard (1947), Stark (1958).

Opisodasys keeni (Baker, 1896) by Allred (1968), Augustson (1941a), Baker (1896, 1904), Coultrip et al. (1973), Easton (1983), Egoscue (1976), Glicken and Schwab (1980), Haas et al. (1980), Hansen (1964), Harvey (1907), Holland (1949), Hubbard (1947, 1949), Jellison (1929), Nelson and Smith (1976), O'Farrell (1975), Stark (1958), Stark and Kinney (1969), Wagner (1936), Zwolak et al. (2013).

Opisodasys nesiotus Augustson, 1941 by Augustson (1941b), Holdenried et al. (1951).

Orchopeas caedens (Jordan, 1925) by Hubbard (1947).

Orchopeas howardi (Baker, 1895) by Baker (1906), Benton et al. (1969), Fox (1940), Frandsen and Grundmann (1960), Geary (1959), Scharf and Stewart (1980), Stark (1958), Stewart (1933).

Orchopeas leucopus (Baker, 1904) by Beck (1966), Benton and Krug (1956), Benton and Altmann (1964), Benton et al. (1971), Brown (1944), Buckner (1964), Connor (1960), Douglas (1969), Egoscue (1976), Elzinga and Rees (1964), Fox (1940), Geary (1959), Haas et al. (1973), Holdenried and Morlan (1956), Holland (1949), Holland and Benton (1968), Hubbard (1947, 1949), Judd (1955), Lawrence et al. (1956), Layne (1958), Main (1970), Morlan (1955), Poorbaugh and Gier (1961), Rapp and Gates (1957), Reeves et al. (2007), Robert (1962), Scharf and Stewart (1980), Scholten et al (1962), Stark (1958), Storm and Ritzi (2008), Verts (1961), Whitaker and Corthum (1967), Whitaker et al. (1975), Wilson (1961), Wyman and Schaefer (1972), Ford et al. (2004).

Orchopeas neotomae Augustson 1943 by Haas et al. (1973), Stark (1958), Ford et al. (2004).

Orchopeas nepos (Rothschild, 1905) by Augustson (1941a), Hubbard (1947).

Orchopeas sexdentatus (Baker, 1904) by Allred (1952), Allred (1968), Egoscue (1976), Hubbard (1947), Morlan (1955), Stark (1958), Ford et al. (2004).

Oropsylla bruner (Baker, 1895) by Benton et al. (1971) as *Opisocrostis bruneri*.

Oropsylla idahoensis (Baker, 1904) by Allred (1952), Hansen (1964), and Stark (1958).

Oropsylla labis (Jordan and Rothschild, 1922) by Allred (1968) as *Opisocrostis labis*.

Oropsylla montana (Baker, 1895) by Allred (1952), Beer et al. (1959), Hubbard (1947), Rutledge et al. (1979), Stark (1958), all cited as *Diamanus montanus*.

Peromyscopsylla catatina (Jordan, 1928) by Buckner and Blasko (1969), Lawrence et al. (1965), Main (1970), Reeves et al. (2007), Robert (1962), Scharf and Stewart (1980).

Peromyscopsylla draco Hopkins, 1951 by Morlan (1955), Ford et al. (2004).

Peromyscopsylla ebrighti (C. Fox, 1926) by Hubbard (1947), Hardy et al. (1974).

Peromyscopsylla hamifer vigens Jordan, 1937 from New Mexico by Haas et al. (1973), Ford et al. (2004); by Egoscue (1976).

Peromyscopsylla hesperomys (Baker, 1904) by Allred (1968), Augustson (1941a), Beck and Allred (1966), Beck (1966), Beer et al. (1959), Benton and Krug (1956), Benton and Altmann (1964), Douglas (1969), Easton (1983), Egoscue (1976), Fox (1940), Geary (1959), Haas et al. (1973), Holdenried et al (1955), Holland (1949), Holland and Benton (1968), Hubbard (1947), Jameson and Brennan (1957), Kinsella (1968), Morlan (1955), Nelson and Smith (1976), Reeves et al. (2007), Scharf and Stewart (1980), Stark (1958), Wyman and Schaefer (1972), Ford et al. (2004), Zwolak et al. (2013). By Stanford (1931), cited as *Ctenopsyllus hesperomys*.

Peromyscopsylla scotti I. Fox, 1939 by Poorbaugh and Gier (1961), Reeves et al. (2007).

Peromyscopsylla selenis (Rothschild, 1906) by Easton (1983), Egoscue (1976), Haas et al. (1973), Holland (1949), Hubbard (1947), Kinsella and Pattie (1967), Rothschild (1906), Stark (1958), Stark and Kinney (1969), Stewart (1933), Wagner (1936), Ford et al. (2004), Zwolak et al. (2013). From Tennessee by Shaftesbury (1934), cited as *Ctenopsyllus selenis*.

Phalacropsylla allos Wagner, 1936 by Ford et al. (2004), Nelson and Smith (1976), Senger (1966), Stark and Kinney (1969).

Phalacropsylla oregonensis Lewis and Maser 1978 by Lewis and Maser (1978).

Phalacropsylla paradisea Rothschild, 1915 by Allred (1968).

Phalacropsylla sp. from Benton County, Washington, by O'Farrell (1975).

Pleochaetis exilis (Jordan, 1937) by Allred (1968), Stark (1958), both cited as *Monopsyllus exilis*.

Plusaetis aztecus Barrera, 1954 by Tipton and Mendez (1968), cited as *Pleochaetis aztecus*.

Plusaetis sibynus (Jordan, 1925) by Beer et al. (1959) and Hubbard (1947), Tipton and Mendez (1968), all cited as *Pleochaetis sibynus*.

Rhadinopsylla fraterna (Baker, 1895) by Holland (1949), Woods and Larson (1971), Zwolak et al. (2013). By Allred (1968) as *Rectofrontia fraterna*.

Rhadinopsylla goodi (Hubbard, 1941) by Elzinga and Rees (1964), Hubbard (1947).

Rhadinopsylla rauschi Holland, 1979 by Holland (1979).

Rhadinopsylla sectilis Jordan and Rothschild, 1923 by Allred (1968), Easton (1983), Egoscue (1966, 1976), Ford et al. (2004), Haas et al. (1973), Holland (1949), Hubbard (1947), Morlan (1955), Senger (1966), Stark (1958), Stark and Kinney (1969), Wagner (1936). Also by Dunn and Parker (1923), cited as *Micropsylla peromyscus*.

Stenistomera alpina (Baker, 1895) by Hubbard (1947), Egoscue (1976), Stark (1958).

Stenistomera hubbardi Egoscue 1968 by Egoscue (1968).

Stenistomera macrodactyla (Good, 1942) by Allred (1968), Beck (1966), Douglas (1969). By Stark (1958), cited as *Miochaeta macrodactyla*.

Stenoponia americana (Baker, 1899) by Eads et al. (1979), Ford et al. (2004), Haas et al. (1973), Holland (1949), Layne (1958), Poorbaugh and Gier (1961), Reeves et al. (2007), Scharf and Stewart (1980), Senger (1966), Whitaker and Corthum (1967), Whitaker et al. (1975).

Stenoponia ponera Traub and Johnson, 1952 by Beer et al. (1959).

Thrassis acamantis (Rothschild, 1905) by Hubbard (1947), Stark (1958).

Thrassis aridis Prince, 1944 by Morlan (1955), Stark (1958).

Thrassis augustsoni Hubbard, 1949 by Hardy et al. (1974).

Thrassis bacchi pansus (Jordan, 1925) by Ford et al. (2004), Morlan (1955), all cited as *Thrassis pansus*.

Thrassis campestris Prince, 1944 by Prince (1944), Ford et al. (2004).

Thrassis fotus (Jordan, 1925) by Schwan and Dobkin (1981).

Thrassis francisi (C. Fox, 1927) by Allred (1968), Stark (1958).

Thrassis pandorae Jellison, 1937 by Stark (1958).

Thrassis pansus (Jordan, 1925) by Ford et al. (2004), Morlan (1955).

Tunga monositus Barnes and Radovsky, 1969 by Barnes and Radovsky (1969), Camargo et al. (2017).

Acanthocephala

Moniliformis clarki in Pine County, Minnesota, by Erickson (1938a,b), reported by Van Cleave (1953); from Utah by Grundman (1959), Grundmann and Frandsen (1959, 1960), Grundman et al. (1961, 1976); Crook (1964), Crook and Grundmann (1964b), including life history; from California by McKeever (1963); from central and southern Rocky Mountains by Smith (1953); from Colorado by Douglas (1969), Leiby (1961), Stock (1962); from Illinois by Barker et al. (1987).

Moniliformis sp., immature, from Nevada by Babero and Matthias (1967).

Nematodes

Ascaris lumbricoides, experimental infections of mice by Tiner (1953).

Aspicularia sp. from Colorado by Stock (1962).

Aspicularis americana from Minnesota by Erickson (1938a,b), Mathies (1958); from Montana by Zwolak et al. (2013).

Aspicularis tetraptera from *P. m. sonoriensis* from the southern arm of Great Salt Lake Desert in Utah by Grundmann (1957a), Grundmann and Frandsen (1960), Mathies (1958).

Baylisascaris procyonis, experimental infections in *P. leucopus*, *P. maniculatus*, *P. californicus*, and *P. polionotus* by Sapp et al. (2016a,b), Sapp et al. (2018). Logiudice (2001), Page (2013), and Page et al. (2001) point out that latrine foraging by two small mice allow transmission of *B. procyonis*.

Brevistriata skrjabini from Utah by Frandsen and Grundmann (1961b).

Carolinensis carolinensis from Hidalgo, Mexico, by Pulido-Flores et al. (2005).

Capillaria americana by Read (1949a,b).

Capillaria gastrica from Hidalgo, Mexico, by Pulido-Flores et al. (2005).

Capillaria hepatica by Freeman (1958), Freeman and Wright (1960), Herman (1981), Lubinsky (1956, 1957), Wright (1961), Lubinsky et al. (1971), Rausch (1961), Layne (1968), Zenchak and Hall (1971); Solomon and Handley (1971). See also Layne and Griffo Jr. (1960), Stock (1962), Meagher (1998, 1999), Dalquest (1948). By Zwolak et al. (2013) cited as *Calodium hepaticum*.

Capillaria sp. from Quebec by Schad (1956); from Utah by Grundmann et al. (1961).

Filariid nematode larvae (*Filaroides martis*, *Perostrongylus pridhami*, or *Trilobostrongylus bioccai*) from Algonquin Park, Ontario, Canada, by Lankester and Anderson (1966).

Gongylonema mysciphilia from Utah by Frandsen and Grundmann (1961a), Grundmann et al. (1961).

Gongylonema peromysci from Nevada by Babero and Matthias (1967); from Arizona by Kruidenier and Peebles (1958).

Gongylonema sp. from Colorado by Hall (1912); from Utah by Kruidenier and Peebles (1957), Frandsen and Grundmann (1961b); from Arizona by Kruidenier and Peebles (1957).

Heligmosomoides polygyrus from California by Ehrenford (1954). Described as *H. p. bakeri* by Durette-Desset et

al. (1972) but found not to be infective by Forrester and Neilson (1973); Forrester (1971) found it not to be infective to *P. maniculatus* after 16 years of passage in laboratory mice. Cable et al. (2006) reported molecular evidence that *H. polygyrus* from lab mice and wood mice are separate species. See also Behnke et al. (1991).

Heligmosomoides vandegrifti from Pennsylvania by Durette-Desset and Kinsella (2007); from Montana by Zwolak et al. (2013).

Longistriata carolinensis from Indiana by Dikmans (1935); from North Carolina by Durette-Desset (1974), who cited it as *Boreostrongylus carolinensis* (Dickmans, 1935) n. comb.

Mastophorus dipodemi from Sevilleta National Wildlife Refuge, New Mexico, unpublished (Arctos Database).

Mastophorus muris from Quebec by Schad (1954, 1956).

Mastophorus numidica from Utah by Crook and Grundmann (1964a); they reported *Eleodes tuberculata patruelis* as natural and experimentally infected intermediate hosts. From Colorado by Dyer and Olsen (1967a,b), who reported grasshoppers (*Melanoplus femurrubrum*), crickets (*Acheta domestica*), and beetles (*Eleodes obsoleta*) as intermediate hosts. Also from Colorado by Douglas (1969).

Nematospiroides dubius from Nevada by Babero and Matthias (1967); from near Davis, California, by Ehrenford (1954); life-cycle studies by Baker (1954), Cross (1960), Ehrenford (1954), and Cross and Duffy (1963); see also Behnke et al (1991); Yamaguti (1958–1963).

Nippostrongylus brasiliensis from Utah by Grundmann and Frandsen (1960); listed as a parasite of *P. maniculatus* by Erickson (1938a,b).

Nippostrongylus muris reported from deer mice by Porter (1935); from Utah by Grundmann and Frandsen (1960), Frandsen and Grundmann (1961b).

Nippostrongylus sp. from Idaho by Leiby (1962). Haley (1961) reported that Porter's (1935) attempt to infect *P. maniculatus* was not successful. Haley (1961) examined the specimens of Grundmann and Frandsen and indicated that they were not *N. brasiliensis*. Travassos and Darriba (1929) reported that *N. muris* (Yokogawa, 1920) was a synonym of *N. brasiliensis*.

Parastrongylus schmidti, experimental infection by Kinsella (1987).

Physaloptera sp. larvae from Colorado by Dyer and Olsen (1967b); from Montana by Zwolak et al. (2013).

Pterygodermatites peromysci from Maryland, Virginia, Wisconsin, Georgia, and Florida by Lichtenfels (1970).

Protospirura numidica from Utah by Grundmann (1957a, 1957b), Healey and Grundmann (1974), Grundmann and Frandsen (1959, 1960), Frandsen and Grundmann (1961b), Grundmann et al. (1976), Grundmann and Warnock (1964), Crook (1964), Crook and Grundmann (1964a); from Colorado by Leiby (1961); from Idaho by Leiby (1962); from Montana by Zwolak et al. (2013). Life-cycle study by Crook and Grundmann (1964a) using *Eleodes tuberculata patrulis* as a natural intermediate host. Dyer and Olsen (1967) experimentally infected a cricket, *Acheta domestica* (L.); beetle, *Eleodes obsoleta* (Say); and grasshopper, *Melanoplus femurrubrum* (DeGeer). Quentin et al. (1968) established infections in the linear earwig, *Doru lineare* (Eschscholtz), and cockroach, *Periplaneta americana* (Linn.). Healey and Grundmann (1974) reported *Eleodes tuberculata*, *Melanoplus femurrubrum* and *M. atlanis*, and *Gryllus pennsylvanicus* as intermediate hosts.

Protospirura peromysci from Nevada by Babero and Matthias (1967); from Chanel Islands, California, by Smith and Carpenter (2006).

Protospirura sp. from Washington by O'Farrell (1975).

Rictularia coloradensis from Arizona by Kruidenier and Peebles (1957); from Colorado by Hall (1916), Stock (1962), Leiby (1961), Dyer and Olsen (1967b); from Colorado and Utah by Lichtenfels (1970); from Utah by Grundmann and Frandsen (1959, 1960), Grundmann et al. (1976), Grundmann and Warnock (1964); from Illinois by Barker et al. (1987); from Nevada by Babero and Matthias (1967); from Washington by Rankin (1945); from Idaho by Leiby (1962); from Wisconsin by Tiner (1948b); from Washington by Rankin (1945). *R. coloradensis* is a common parasite in northern New Mexico.

Oswald (1958a,b), working with specimens obtained from *P. leucopus*, found that the larvae develop to the infective stage in Orthoptera, wood roaches (*Parcoblatta* spp.), and camel crickets (*Ceuthophilus* spp.), and all were natural intermediate hosts. He reported that in natural infections of *Peromyscus* 76.5% harbored *R. coloradensis*.

Rictularia onychomis from Nevada by Babero and Matthias (1967).

Rictularia sp. from Utah by Grundmann (1957a); from Washington by O'Farrell (1975).

Strongylida lungworms. Encapsulated larvae in lungs from Algonquin Park, Ontario, Canada, by Lankester and Anderson (1966).

Syphacia obvelata from Washington by Rankin (1945); from Colorado by Douglas (1969).

Syphacia peromysci from Grand Canyon, Arizona; Sevilleta, New Mexico, unpublished (Arctos Database); from Wisconsin in *P. maniculatus bairdii* by Tiner (1948a), Tiner and Rausch (1950); from Quebec, Canada, by Schad (1956); from Utah by Grundmann (1957b, 1959), Grundmann and Frandsen (1959, 1960), Grundmann and Warnock (1964), Grundmann et al. (1976), Frandsen

and Grundmann (1961b), Kruidenier and Peebles (1957), Ogden (1971), Quentin and Kinsella (1972); from the Navajo Reservoir Basin, Colorado, and New Mexico by Stock (1961); from Arizona by Kruidenier and Peebles (1957); from Idaho by Leiby (1962); from Montana by Zwolak et al. (2013); from Hidalgo, Mexico, by Pulido-Flores et al. (2005); from Arkansas by McAllister et al. (2020).

Syphacia samorodini from Stanbury Mountains, Utah, by Grundmann (1957a); from Minnesota by Erickson (1938b); see also Kruidenier et al. (1961).

Syphacia spp. larvae from Lubbock County, Texas, by Rodenberg and Pence (1978).

Trichinella spiralis. None found in 472 animals from Mountain Lake, Virginia, by Solomon and Warner (1969). Experimental infections of *T. nativa* and *T. pseudospiralis* by Poirier (1994), Poirier at al. (1993). No infections from New Jersey by Leiby et al. (1988).

Trichuris muris from deer mice with prepatent period of 5 weeks by Worley et al. (1960); in the Channel Islands, California, by Smith and Carpenter (2006).

Trichuris perognatha from Utah by Grundmann and Frandsen (1960); from Colorado by Dyer and Olsen (1967b).

Trichuris sp. from Utah by Frandsen and Grundmann (1961a), Grundmann et al. (1961).

Trichuris stansburyi from Utah by Frandsen and Grundmann (1961a); from Colorado by Stock (1962).

Wellcomia sp. from Washington by O'Farrell (1975).

Trematodes

Brachylaima (*Postharmostomum*) *sexconvolutum* from Ohio by Miller (1936).

Brachylaima sp. from Utah by Grundmann and Frandsen, (1959).

Brachylaime microti from Utah by Frandsen and Grundmann (1960, 1961b), Grundmann and Frandsen (1960); from Texas by Redetzke and Canaris (1977), Bristol and Canaris (1973).

Concinnum peromysci by Neidert and Macy (1968).

Euryhelmis pacificus from the northwest US by Senger and Macy (1952).

Metacercariae from Quebec, Canada, by Schad, (1956).

Postharmostomum helicis from Quebec, Canada, by Schad (1956); experimental infection by Robinson (1949), Ulmer (1951).

Postharmostomum sexconvolutum listed by Yamaguti (1958–1963).

Quinqueserialis quinqueserialis could not be established in *P. maniculatus* by experimental infections, Kinsella (1971).

Schistosomatium douthitti, experimental infections by Short (1951, 1952); Schwanz (2006), Price (1931), Malek (1977).

Cestodes

Anoplocephala sp. from Colorado by Hall (1912).

Catenotaenia dendritica (*C. peromysci* according to Wolfgang (1956) from Quebec, Canada, Schad (1954); from Colorado by Leiby (1961); from Idaho by Leiby (1962).

Catenotaenia linsdaeli from Utah by Grundmann (1958), Grundmann and Frandsen (1959, 1960), Frandsen and Grundmann (1961b).

Catenotaenia peromysci from Coyote, Rio Arriba County, New Mexico, and from Colorado and Wyoming by Smith (1954); from Alberta, Canada, by Lubinsky (1957); from California by Voge (1955); from Utah by Grundmann et al. (1976), Bienek and Grundmann (1974); from Arkansas by McAllister et al. (2020). Cited by Hwang et al (2007), Hansen (1950).

Choanotaenia peromysci by Erickson (1938a,b), Hansen (1950) (syn. *Prochoanotaenia peromysci* Erickson 1938); from Alberta by Lubinsky (1957); from Colorado by Leiby (1961); from California by Voge (1955); from the southern Rocky Mountains by Smith (1954); from South Dakota by Ulrich and Vaughn (1963); from Illinois by Barker et al. (1987); from Montana by Zwolak et al. (2013).

Cladotaenia circi larvae from Utah by Grundmann and Frandsen (1959, 1960), Frandsen and Grundmann (1961b).

Cladotaenia globifera larva from Algonquin Park, Ontario, Canada, by Freeman and Wright (1960).

Cladotaenia sp. from Minnesota by Erickson (1938b).

Cysticercus sp. Skinker (1935).

Echinococcus multilocularis hydatid cyst from Minnesota by Carney and Leiby (1968); from North Dakota by Leiby (1965) and S. H. Richards, North Dakota Game and Fish Department, Bismarck (unpublished data) reported by Rausch (1967), Leiby et al. (1970), Leiby and Kritsky (1974). According to Jones and Pybus (2001), *P. maniculatus* is one of two main intermediate hosts for *E. multilocularis* in North America. Leiby and Kritsky (1974), Kritsky and Leiby (1975), from Minnesota and Winnipeg, Manitoba, Canada, by Leiby et al. (1969), Lee (1969); from Manitoba by Lubinsky et al. (1971); from Montana by Feigley and Worley (1988); from Alberta, Canada, by Holmes et al. (1971), Chalmers and Barrett (1974); from New Mexico by Botero-Canola et al. (2019). Rausch and Richards (1971) feeding experiments show that meadow voles are more susceptible than deer mice. Summary of life cycles in north-central US by Leiby and Kritsky (1972). Ecology and distribution in Wells County, North Dakota, by Ohbayashi et al. (1971). Jacobsen (1968) reported no infections in *P. maniculatus* from grain fields around the Winnipeg, Manitoba, vicinity, but it was present around the Truax Mine in the same area by Leiby et al. (1969).

Echinococcus sibiricensis from St. Lawrence Island, Alaska, by Rausch and Schiller (1954), who indicate a single specimen of *Peromyscus* sp. was infected, but rodents of this genus appear to be relatively refractory (Rausch and Schiller, 1954). Subsequent work shows a new biotype of *E. multilocularis* may be in the lower 48 states (Botero et al., 2019).

Hydatigera lyncis by Skinker (1935) from Virginia, by Holloway (1963).

Hymenolepis bennetti from Ontario by Freeman (1960); from Illinois by Barker et al. (1987).

Hymenolepis citelli from Utah by Crook (1964), Grundmann and Frandsen (1959, 1960); Frandsen and Grundmann (1961b), Grundmann et al. (1976), Cook (1964); some host resistance was reported by Wassom et al. (1973, 1974, 1986).

Hymenolepis diminuta from Northrup Canyon, Washington, by Rankin (1945).

Hymenolepis fraterna, experimental infection by Hunninen (1935).

Hymenolepis peromysci by Tinkle (1972) in *P. boylii* and also in *P. maniculatus* from California. Tinkle reported that stink beetles *Eleodes armata* LeConte and *E. dentipes* Eschscholtz were intermediate hosts. Growth and development in hamsters by Stallard and Arai (1978).

Hymenolepis sp. cysticercoid from Idaho by Leiby (1962); from Nevada by Babero and Matthias (1967).

Hymenolepis sp. from West Virginia by Zenchak and Hall (1971); from Nevada by Babero and Matthias (1967); from Minnesota by Erickson (1938a,b), Hansen (1950); from Nebraska by Hansen (1950); from Montana by Zwolak et al. (2013).

Mesocestoides carnivoricolus tetrathyridia from Utah by Grundmann (1956, 1957a, 1958), Grundmann and Frandsen (1959, 1960); Frandsen and Grundmann (1961b).

Mesocestoides tetrathyridia from Taos County, New Mexico, by Ubelaker et al. (2014); from Coos County, Oregon, by Kegley et al. (1970); from Barkley Sound, British Columbia, by Herman (1981).

Paranoplocephala sp. from Utah by Grundmann and Frandsen (1959, 1960).

Paruterina candelabraria plerocercoid from Canada by Baron (1971); from Utah by Grundmann and Frandsen (1959); Frandsen and Grundmann (1961b); from Wisconsin with life cycle by Rausch (1949).

Paruterina rauschi plerocercoid in liver from West Virginia by Zenchak and Hall (1971); from Algonquin Park, Ontario, Canada, by Freeman and Wright (1960).

Prochoanotaenia sp. by Erickson (1938a, b).

Taenia mustelae cysticercus in liver and lungs from West Virginia by Zenchak and Hall (1971); from Ontario by Freeman (1956); from Algonquin Park, Ontario, Canada, by Freeman and Wright (1960).

Taenia ovis from Quebec and the coast of Labrador by Schad (1954).

Taenia rileyi by Riser (1956).

Taenia sp. from Hidalgo, Mexico, by Pulido-Flores et al. (2005).

Taenia taeniaeformis from California by Theis and Schwab (1992).

Taenia tenuicollis (= *T. hydatigena*) cysticercus from Montana and Oregon by Locker (1955); from Quebec by Schad (1956).

Other parasites carried by **Peromyscus**

Clostridium botulinum, which results in botulism. Botulism type 3 is found in aquatic birds and is known to infect mice in laboratory conditions.

Mice, including *Peromyscus* spp., can also serve as carriers of parasitic organisms of other animals. Davidson and Nettles (1988) report that mice can serve to carry these parasites from other animals.

Pseudorabies Aujeszky's disease caused by a herpesvirus in swine and cattle, sheep, horses, dogs, cats, foxes, raccoons, skunks, rats, and mice.

Stomach worms *Gnathostoma procyonis* and *Physaloptera rara* and larvae from crickets can be stored in mice and later transmitted to raccoons when they prey on mice.

Toxoplasma gondii, which results in toxoplasmosis. Experimental infections in mice suggest mice can carry the protozoan and allow transmission to cats or other wild or domestic felines.

Literature cited

Adams, W.; Emmons, R.; and Brooks, J. 1970. The changing ecology of murine (endemic) typhus in Southern California. The American Journal of Tropical Medicine and Hygiene 19: 311–318.

Adjemian, J. Z.; Adjemian, M. K.; Foley, P.; Chomel, B. B.; Kasten, R. W.; and Foley, J. E. 2008. Evidence of multiple zoonotic agents in a wild rodent community in the eastern Sierra Nevada. Journal of Wildlife Diseases 44: 737–742. https://doi.org/10.7589/0090-3558-44.3.737

Allred, D. M. 1952. Plague important fleas and mammals in Utah and the western United States. The Great Basin Naturalist 12: 67–75.

Allred, D. M. 1957a. Mites found on mice of the genus *Peromyscus* in Utah. II. Family Haemogamasidae. Proceedings of the Entomoligcal Society of Washington 59: 31–39.

Allred, D. M. 1957b. Mites found on mice of the genus *Peromyscus* in Utah. III. Family Dermanyssidae. The American Midland Naturalist 57: 450–460. https://doi.org/10.2307/2422410

Allred, D. M. 1957c. Mites found on mice of the genus *Peromyscus* in Utah. V. Trombiculidae and miscellaneous families. The Great Basin Naturalist 17: 95–102.

Allred, D. M. 1958. Mites found on mice of the genus *Peromyscus* in Utah. IV. Families Laelaptidae and Phytoseiidae. Pan-Pacific Entomologist 34: 17–32.

Allred, D. M.; Beck, D. E.; and White, L. D. 1960. Ticks of the genus *Ixodes* in Utah. Brigham Young University Science Bulletin, Biological Series 1: 1–42.

Allred, D. M.; and Goates, M. A. 1964. Mites from mammals at the Nevada test site. The Great Basin Naturalist 24: 71–73.

Allred, D. M. 1965. Clarification of type data for *Ischyropoda furmani* Keegan. The Journal of Parasitology 51: 604.

Allred, D. M.; and Beck, D. E. 1966. Mites of Utah mammals. Brigham Young University Science Bulletin, Biological Series 8: 1–123.

Allred, D. M. 1968. Fleas of the national reactor testing station. Great Basin Naturalist 28: 73–87.

Allred, D. M. 1970. Mites and lice of the National Reactor Testing Station. Brigham Young University Science Bulletin, Biological Series 12: 1–17.

Amman, B. R.; Manangan, A. P.; Flietstra, T. D.; Calisher, C. H.; Carroll, D. S.; Wagoner, K. D.; and Mills, J. N. 2013. Association between movement and Sin Nombre virus (Bunyaviridae: Hantavirus) infection in North American deermice (*Peromyscus maniculatus*) in Colorado. Journal of Wildlife Diseases 49: 132–142. https://doi.org/10.7589/2012-02-041

Augustson, G. F. 1941a. Ectoparasite-host records from the Sierran region of east-central California. Bulletin of the Southern California Academy of Sciences 40: 147–157.

Augustson, G. F. 1941b. Contributions from the Los Angeles Museum–Channel Islands Biological Survey. No. 20. Three new fleas (Siphonaptera). Bulletin of the Southern California Academy of Sciences 40: 101–107.

Babero, B.B.; and Matthias, D. 1967. *Protospirura peromysci* n. sp. (Nematoda: Spiruridae) and other helminths from *Peromyscus* spp. in Nevada. Proceedings of the Helminthological Society of Washington 34: 255–261.

Bacon, M.; and Drake, C. H. 1958. Comparative susceptibility of various species of mice native to Washington to inoculation with virulent strains of *Pasteurella pestis*. The Journal of Infectious Diseases 102: 14–22.

Bacon, M.; Drake, C. H.; and Miller, N. G. 1959. Ticks (Acarina: Ixodoidea) on rabbits and rodents of eastern and central Washington. The Journal of Parasitology 45: 281–286.

Bai, Y.; Calisher, C. H.; Kosoy, M. Y.; Root, J. J.; and Doty, J. B. 2011. Persistent infection or successive reinfection of deer mice with *Bartonella vinsonii* subsp. *arupensis*. Applied and Environmental Microbiology 77: 1728–1731. https://doi.org/10.1128/AEM.02203-10

Baker, C. F. 1896. A new *Pulex* from Queen Charlotte Islands. The Canadian Entomologist 28: 234–234.

Baker, C. F. 1904. A revision of American Siphonaptera, or fleas; together with a complete list and bibliography of the group. Proceedings of the United States National Museum 27: 365–469.

Baker, C. F. 1905. The classification of the American Siphonaptera. Proceedings of the United States National Museum 29: 121–170.

Baker, N. F. 1954. Trichostrongylidosis—the mouse as an experimental animal. Proceedings of the American Veterinary Medical Association. 91st Annual Meeting. Seattle, pp. 185–191.

Bakerspigel, A. 1956. *Haplosporangium* in Saskatchewan rodents. Mycologia 48: 568–572.

Barker, C.; Dyer, W.; and Feldhamer, G. 1987. Helminths of *Peromyscus leucopus*, *P. maniculatus*, and *Blarina carolinensis* from Southern Illinois. Transactions of the Illinois Academy of Science 80: 119–127.

Barnes, A. M.; and Radovsky, F. J. 1969. A new *Tunga* (Siphonaptera) from the Nearctic Region with description of all stages. Journal of Medical Entomology 6: 19–36. https://doi.org/10.1093/jmedent/6.1.19

Barnes, A. M.; Tipton, V. J.; and Wildie, J. A. 1977. The subfamily Anomiopsyllinae (Hystrichopsyllidae: Siphonaptera). I. A revision of the genus *Anomiopsyllus* Baker. The Great Basin Naturalist 37: 138–206.

Barnes, A. M. 1982. Surveillance and control of bubonic plague in the United States. Symposium of the Zoological Society of London 50: 237–268.

Baron, R. W. 1971. The occurrence of *Paruterina candelabraria* (Goeze, 1782) and *Cladotaenia globifera* (Batsch, 1786) in Manitoba. Canadian Journal of Zoology 49: 1399–1400. https://doi.org/10.1139/z71-208

Bartlow, A. W. 2019. Histological findings in wild rodents of the Great Basin. Western North American Naturalist 79: 323–336.

Beck, D. E. 1955. Some unusual distribution records of ticks in Utah. The Journal of Parasitology 41: 1–4.

Beck, D. E.; Allred, D. M.; and Brinton, E. P. 1963. Ticks of the Nevada test site. Brigham Young University Science Bulletin, Biological Series 4: 1–11.

Beck, D. E. 1966. Siphonaptera (fleas) of Mesa Verde National Park, Montezuma, Colorado. Great Basin Naturalist 26: 76–78.

Beck, D. E.; and Allred, D. M. 1966. Siphonaptera (fleas) of the Nevada test site. Brigham Young University Science Bulletin, Biological Series 7: 1–27.

Beer, J. R.; Cook, E. F.; and Schwab, R. G. 1959. The ectoparasites of some mammals from the Chiricahua Mountains, Arizona. The Journal of Parasitology 45: 605–613.

Behnke, J. M.; Keymer, A. E.; and Lewis, J. W. 1991. *Heligmosomoides polygyrus* or *Nematospiroides dubius*? Parasitology Today 7: 177–179.

Bennett, S. G.; Webb Jr., J. P.; Madon, M. B.; Childs, J. E.; Ksiazek, T. G.; Torrez-Martinez, N.; and Hjelle, B. 1999. Hantavirus (Bunyaviridae) infections in rodents from Orange and San Diego counties, California. The American Journal of Tropical Medicine and Hygiene 60: 75–84. https://doi.org/10.4269/ajtmh.1999.60.75

Bennett, S. G.; Milazzo, M. L.; Webb Jr., J. P.; and Fulhorst, C. F. 2000. Arenavirus antibody in rodents indigenous to coastal southern California. The American Journal of Tropical Medicine and Hygiene 62: 626–630. https://doi.org/10.4269/ajtmh.2000.62.626

Bennett, S. G.; Comer, J. A.; Smith, H. M.; and Webb, J. P. 2007. Serologic evidence of a *Rickettsia akari*–like infection among wild-caught rodents in Orange County and humans in Los Angeles County, California. Journal of Vector Ecology 32: 198–201. https://doi.org/10.3376/1081-1710(2007)32[198:SEOARA]2.0.CO;2

Benton, A. H.; and Krug, R. F. 1956. Mammals and siphonapterous parasites of Rensselaer County, New York. New York State Museum and Science Service Bulletin. University of the State of New York.

Benton, A. H.; and Altmann, H. J. 1964. A study of fleas found on *Peromyscus* in New York. Journal of Mammalogy 45: 31–36. https://doi.org/10.2307/1377291

Benton, A. H.; Tucker, H. H.; and Kelly, D. L. 1969. Siphonaptera from Northern New York. Journal of the New York Entomological Society 77: 193–198.

Benton, A. H.; Larson, O. R.; and Ven-Huizen, B. A. 1971. Siphonaptera from Itasca State Park region. Journal of the Minnesota Academy of Science 37: 91–92.

Bienek, G. K.; and Grundmann, A. W. 1974. *Catenotaenia utahensis* sp. n. (Cestoda: Catenotaeniidae) from Merriam kangaroo rat, *Dipodomys merriami vulcani*, in Utah. Proceedings of the Helminthological Society of Washington 41: 134–139.

Bishop, F. C.; and Trembley, H. L. 1945. Distribution and hosts of certain North American ticks. The Journal of Parasitology 31: 1–54.

Blair, W. F. 1942. Size of home range and notes on the life history of the woodland deer-mouse and eastern chipmunk in northern Michigan. Journal of Mammalogy 23: 27–36.

Bledsoe, B. 1979. Sporogony of *Sarcocystis idahoensis* in the gopher snake, *Pituophis melanoleucus* (Daudin). The Journal of Parasitology 65: 875–879. https://doi.org/10.2307/3280240

Boeck, W. G.; and Stiles, C. W. 1923. Studies on various internal parasites (especially Amoebae) of man. United States Public Health Service. Hygienic Laboratory Bulletin No 133: 1–202.

Boone, J. D.; McGwire, K. C.; Otteson, E. W.; DeBaca, R. S.; Kuhn, E. A.; Villard, P.; Brussard, P. F.; and St. Jeor, S. C. 2000. Remote sensing and geographic information systems: charting Sin Nombre virus infections in deer mice. Emerging Infectious Diseases 6: 248–258.

Botero-Cañola, S.; Dursahinhan, A. T.; Rácz, S. E.; Lowe, P. V.; Ubelaker, J. E.; and Gardner, S. L. 2019. The ecological niche of *Echinococcus multilocularis* in North America: understanding biotic and abiotic determinants of parasite distribution with new records in New Mexico and Maryland, United States. Therya 10: 91–102. https://doi.org/10.12933/therya-19-749

Bowen, G. S.; Shriner, R. B.; Pokorny, K. S.; Kirk, L. J.; and McLean, R. G. 1981a. Experimental Colorado tick fever virus infection in Colorado mammals. The American Journal of Tropical Medicine and Hygiene 30: 224–229. https://doi.org/10.4269/ajtmh.1981.30.224

Bowen, G. S.; McLean, R. G.; Shriner, R. B.; Francy, D. B.; Pokorny, K. S.; Trimble, J. M.; Bolin, R. A.; Barnes, A. M.; Calisher, C. H.; and Muth, D. J. 1981b. The ecology of Colorado tick fever in Rocky Mountain National Park in 1974. The American Journal of Tropical Medicine and Hygiene 30: 490–496. https://doi.org/10.4269/ajtmh.1981.30.490

Bowers, J. H.; Hayes, R. O.; and Hughes, T. B. 1969. Studies on the role of mammals in the natural history of Western Encephalitis in Hale County, Texas. Journal of Medical Entomology 6: 175–178. https://doi.org/10.1093/jmedent/6.2.175

Brennan, J. M. 1946. A new genus and species of chigger, *Chatia setosa* (Trombiculidae, Acarina) from northwestern United States. The Journal of Parasitology 32: 132–135.

Brennan, J. M. 1948. New North American chiggers (Acarina; Trombiculidae). The Journal of Parasitology 34: 465–478.

Brennan, J. M. 1965. Two new species and other records of chiggers from Texas (Acarina: Trombiculidae). Acarologia 7: 79–83.

Brennan, J. M.; and D. E. Beck 1955. The chiggers of Utah (Acarina: Trombiculidae). Great Basin Naturalist 15: 1–26.

Brennan, J. M.; and Jones, E. K. 1954. A report on the chiggers (Acarina: Trombiculidae) of the Frances Simes Hastings Natural History Reservation, Monterey County, California. The Wasmann Journal of Biology 12: 155–194.

Brennan, J. M.; and Wharton, G. W. 1950. Studies on North American chiggers. No. 3. The subgenus *Neotrombicula*. The American Midland Naturalist 44: 153–197.

Bristol J. R.; and Canaris, A. G. 1973. Physiology, ecology and life cycle of *Brachylaime microti* (Trematoda). American Philosophical Society Year Book, 1973, 296–298.

Brown, J. H. 1944. The fleas (Siphonaptera) of Alberta, with a list of the known vectors of sylvatic plague. Annals of the Entomological Society of America 37: 207–213.

Brown, J.H.; and J. M. Brennan 1952. A note on the chiggers (Trombiculidae) of Alberta. Canadian Journal of Zoology 30: 338–343.

Buchholz, M. J.; Davis, C.; Rowland, N. S.; and Dick, C. W. 2018. *Borrelia burgdorferi* in small mammal reservoirs in Kentucky, a traditionally non-endemic state for Lyme disease. Parasitology Research 117: 1159–1167. https://doi.org/10.1007/s00436-018-5794-x

Buckner, C. H. 1964. Fleas (Siphonaptera) of Manatoba mammals. The Canadian Entomologist 96: 850–856.

Buckner, C. H.; and Blasko, G. G. 1969. Additional range and host records of the fleas (Siphonaptera) of Manitoba. Manitoba Entomologist 3: 65–69.

Burgdorfer, W. 1960. Colorado tick fever: II. The behavior of Colorado tick fever virus in rodents. The Journal of Infectious Diseases 107: 384–388.

Burgdorfer, W.; and Eklund, C. M. 1960. Colorado tick fever: I. Further ecological studies in Western Montana. The Journal of Infectious Diseases 107: 379–383.

Burkot, T. R.; Clover, J. R.; Happ, C. M.; DeBess, E.; and Maupin, G. O. 1999. Isolation of *Borrelia burgdorferi* from *Neotoma fuscipes*, *Peromyscus maniculatus*, *Peromyscus boylii*, and *Ixodes pacificus* in Oregon. The American Journal of Tropical Medicine and Hygiene 60: 453–457. https://doi.org/10.4269/ajtmh.1999.60.453

Burkot, T. R.; Schneider, B. S.; Pieniazek, N. J.; Happ, C. M.; Rutherford, J. S.; Slemenda, S. B.; Hoffmeister, E.; Maupin, G. O.; and Zeidner, N. S. 2000. *Babesia microti* and *Borrelia bissettii* transmission by *Ixodes spinipalpis* ticks among prairie voles, *Microtus ochrogaster*, in Colorado. Parasitology 121: 595–599. https://doi.org/10.1017/S0031182000006788

Burkot, T. R.; Maupin, G. O.; Schneider, B. S.; Denatale, C.; Happ, C. M.; Rutherford, J. S.; and Zeidner, N. S. 2001. Use of a sentinel host system to study the questing behavior of *Ixodes spinipalpis* and its role in the transmission of *Borrelia bissettii*, human granulocytic ehrlichiosis, and *Babesia microti*. The American Journal of Tropical Medicine and Hygiene 65: 293–299. https://doi.org/10.4269/ajtmh.2001.65.293

Burroughs, A. L.; Holdenried, R.; Longanecker, D. S.; and Meyer, K. F. 1945. A field study of latent tularemia in rodents with a list of all known naturally infected vertebrates. The Journal of Infectious Diseases 76: 115–119.

Burt, W. H. 1940. Territorial behavior and populations of some small mammals in southern Michigan. Miscellaneous Publications Museum of Zoology, University of Michigan 45: 1–58.

Cable, J.; Harris, P. D.; Lewis, J. W.; and Behnke, J. M. 2006. Molecular evidence that *Heligmosomoides polygyrus* from laboratory mice and wood mice are separate species. Parasitology 133: 111–122.

Calisher, C. H.; Sweeney, W.; Mills, J. N.; and Beaty, B. J. 1999. Natural history of Sin Nombre virus in western Colorado. Emerging Infectious Diseases 5: 126–134. https://doi.org/10.3201/eid0501.990115

Camargo, I.; Romero-Callejas, E.; Cornejo-Latorre, C.; Rios, E.; and Álvarez-Castañeda, S. T. 2017. Prevalence and intensity of flea *Tunga monositus* (Siphonaptera) in an insular population of *Peromyscus maniculatus* (Rodentia) from Northwest Mexico. Mammalia 81: 429–432. https://doi.org/10.1515/mammalia-2016-0013

Campbell, A.; and MacKay, P. R. 1979. Distribution of the American dog tick, *Dermacentor variabilis* (Say), and its small-mammal hosts in relation to vegetation types in a study area in Nova Scotia. Canadian Journal of Zoology 57: 1950–1959. https://doi.org/10.1139/z79-258

Campos, E. G.; and Stark, H. E. 1979. A revaluation of the *Hystrichopsylla occidentalis* group, with description of a new subspecies (Siphonaptera: Hystrichopsyllidae). Journal of Medical Entomology 15: 431–444. https://doi.org/10.1093/jmedent/15.5-6.431

Carey, A. B.; McLean, R. G.; and Maupin, G. O. 1980. The structure of a Colorado tick fever ecosystem. Ecological Monographs 50: 131–151.

Carmichael, J. W. 1961. Fungi from Alberta rodents. Mycopathologia et Mycologia Applicata 14: 129–135.

Carney, W. P.; and Lieby, P. D. 1968. *Echinococcus multilocularis* in *Peromyscus maniculatus* and *Vulpes vulpes* from Minnesota. The Journal of Parasitology 54: 714.

Carver, S.; Trueax, J. T.; Douglass, R.; and Kuenzi, A. 2011. Delayed density-dependent prevalence of sin nombre virus infection in deer mice (*Peromyscus maniculatus*) in central and western Montana. Journal of Wildlife Diseases 47: 56–63. https://doi.org/10.7589/0090-3558-47.1.56

Catalán-Dibene, J.; Johnson, S. M.; Eaton, R.; Romero-Olivares, A. L.; Baptista-Rosas, R. C.; Pappagianis, D.; and Riquelme, M. 2014. Detection of coccidioidal antibodies in serum of a small rodent community in Baja California, Mexico. Fungal Biology 118: 330–339. https://doi.org/10.1016/j.funbio.2014.01.006

Chalmers, G. A.; and Barrett, M. W. 1974. *Echinococcus multilocularis* Leuckart, 1863 in rodents in southern Alberta. Canadian Journal of Zoology 52: 1091–1091. https://doi.org/10.1139/z74-145

Chapin, E. A. 1919. New Species of North American Siphonaptera. Bulletin of the Brooklyn Entomological Society 14.

Childs, J. E.; Ksiazek, T. G.; Spiropoulou, C. F.; Krebs, J. W.; Morzunov, S.; Maupin, G. O.; Gage, K. L.; Rollin, P. E.; Sarisky, J.; Enscore, R. E.; Frey, J. K.; Peters, C. J.; and Nichol, S. T. 1994. Serologic and genetic identification of *Peromyscus maniculatus* as the primary rodent reservoir for a new hantavirus in the southwestern United States. The Journal of Infectious Diseases 169: 1271–1280. https://doi.org/10.1093/infdis/169.6.1271

Clark, G. M.; Clifford, C. M.; Fadness, L. V.; and Jones, E. K. 1970. Contributions to the ecology of Colorado tick fever virus. Journal of Medical Entomology 7: 189–197. https://doi.org/10.1093/jmedent/7.2.189

Clay, C. A.; Lehmer, E. M.; Jeor, S. S.; and Dearing, M. D. 2009. Sin Nombre virus and rodent species diversity: a test of the dilution and amplification hypotheses. PLOS ONE 4: e6467. https://doi.org/10.1371/journal.pone.0006467

Coady, N. R.; and Nickol, B. B. 2000. Assessment of parenteral *Plagiorhynchus cylindraceus* (Acanthocephala) infections in shrews. Comparative Parasitology 67: 32–39.

Connior, M. B.; Durden, L. A.; McAllister, C. T.; Seville, R. S.; Bursey, C. R. and Robison, H. W. 2017. New records of parasites (Apicomplexa, Nematoda, Acari, Anoplura) from rodents in Arkansas. Journal of the Arkansas Academy of Science 71: 211–214.

Connor, P. F. 1960. The small mammals of Otsego and Schoharie Counties, New York. New York State Museum Bulletin. University of the State of New York, State Education Department, Albany, New York.

Cook, E. F.; and Beer, J. R. 1959. The immature stages of the genus *Hoplopteura* (Anoplura: Hoplopteuridae) in North America, with descriptions of two new species. The Journal of Parasititology 45: 405–416.

Cooley, R. A.; and Kohls, G. M. 1941. Three new species of *Ornithodoros* (Acarina: Ixodoidea). Public Health Reports 56: 587–594.

Cooley, R. A.; and Kohls, G. M. 1945. The genus *Ixodes* in North America. National Institute of Health Bulletin No. 184. National Institute of Health, Hamilton, Montana.

Coultrip, R. L.; Emmons, R. W.; Legters, L. J.; Marshall Jr., J. D.; and Murray, K. F. 1973. Survey for the arthropod vectors and mammalian hosts of Rocky Mountain Spotted Fever and plague at Fort Ord, California. Journal of Medical Entomology 10: 303–309. https://doi.org/10.1093/jmedent/10.3.303

Crook, J. R. 1964. The role of intermediate hosts in the ecological distribution of some endoparasites in the Bonneville Basin, Utah. Dissertation Abstracts 25: 2109.

Crook, J. R.; and Grundmann, W. 1964a. The life history of *Protospirura numidica* Seurat, 1914 (Nematoda: Spiruroidea). Proceedings of the Helminthological Society of Washington 31: 255–229.

Crook, J. R.; and Grundmann, W. 1964b. The life history and larval development of *Moniliformis clarki* (Ward, 1917). The Journal of Parasitology 50: 689–693.

Cross, J. H. 1960. The natural resistance of the white rat to *Nematospiroides dubius* and the effect of cortisone on this resistance. The Journal of Parasitology 46: 175–185.

Cross, J. H.; and Duffy, C. E. 1963. *Nematospiroides dubius* in the abnormal host. Annals of the New York Academy of Sciences 113: 88–99. https://doi.org/10.1111/j.1749-6632.1963.tb40660.x

Crossley Jr., D. A.; and Lipovsky, L. J. 1954. Two new chiggers from the central states (Acarina, Trombiculidae). Proceedings of the Entomological Society of Washington 56: 240–246.

Dalquest, W. W. 1948. Mammals of Washington. University of Kansas, Museum of Natural History, Lawrence.

Dalquest, W. W. 1965. New Pleistocene formation and local fauna from Hardeman County, Texas. Journal of Paleontology 39: 63–79. Cited by Hibbard (1968).

Danforth, M. E.; Messenger, S.; Buttke, D.; Weinburke, M.; Carroll, G.; Hacker, G.; Niemela, M.; Andrews, E. S.; Jackson, B. T.; Kramer, V.; and Novak, M. 2020. Long-term rodent surveillance after outbreak of hantavirus infection, Yosemite National Park, California, USA, 2012. Emerging Infectious Diseases 26: 560–567. https://doi.org/10.3201/eid2603.191307

Danforth, M.; Tucker, J.; and Novak, M. 2018. The deer mouse (*Peromyscus maniculatus*) as an enzootic reservoir of plague in California. EcoHealth 15: 566–576. https://doi.org/10.1007/s10393-018-1337-2

Davidson, W. R.; and Nettles, V. F. 1988. Field manual of wildlife diseases in the southeastern United States. Southeastern Cooperative Wildlife Disease Study, Deptment of Parasitology, College of Veterinary Medicine, University of Georgia, Athens, Georgia.

Davis, B. S. 1952. Studies on the trypanosomes of some California mammals. University of California Publications in Zoology 57: 145–250.

Davis, B. S. 1967. *Isospora peromysci* n. sp., *I. californica* n. sp., and *I. hastingsi* n. sp. (Protozoa: Eimeriidae) from four sympatric species of white-footed mice (*Peromyscus*) in Central California. The Journal of Protozoology 14: 575–585. https://doi.org/10.1111/j.1550-7408.1967.tb02044.x

Davis, J. W.; Hardy, J. L.; and Reeves, W. C. 1974. Modoc viral infections in the deer mouse *Peromyscus maniculatus*. Infection and Immunity 10: 1362–1369.

Davis, R. M.; Smith, R. T.; Madon, M. B.; and Sitko-Cleugh, E. 2002. Flea, rodent, and plague ecology at Chuchupate Campground, Ventura County, California. Journal of Vector Ecology 27: 107–127.

Deardorff, E. R.; Nofchissey, R. A.; Cook, J. A.; Hope, A. G.; Tsvetkova, A.; Talbot, S. L.; and Ebel, G. D. 2013. Powassan virus in mammals, Alaska and New Mexico, USA, and Russia, 2004–2007. Emerging Infectious Diseases 19: 2012–2016. https://doi.org/10.3201/eid1912.130319

del Rocío Reyes-Montes, M.; Pérez-Huitrón, M. A.; Ocaña-Monroy, J. L.; Frías-De-León, M. G.; Martínez-Herrera, E.; Arenas, R.; and Duarte-Escalante, E. 2016. The habitat of *Coccidioides* spp. and the role of animals as reservoirs and disseminators in nature. BMC Infectious Diseases 16: 550. https://doi.org/10.1186/s12879-016-1902-7

DeNatale, C. E.; Burkot, T. R.; Schneider, B. S.; and Zeidner, N. S. 2002. Novel potential reservoirs for *Borrelia* sp. and the agent of human granulocytic ehrlichiosis in Colorado. Journal of Wildlife Diseases 38: 478–482. https://doi.org/10.7589/0090-3558-38.2.478

Dikmans, G. 1935. New nematodes of the genus *Longistriata* in rodents. Journal of the Washington Academy of Sciences 25: 72–81.

Dizney, L.; Jones, P. D.; and Ruedas, L. A. 2010. Natural history of Sin Nombre virus infection in deer mice in urban parks in Oregon. Journal of Wildlife Diseases 46: 433–441. https://doi.org/10.7589/0090-3558-46.2.433

Dodds, D. G.; Martell, A. M.; and Yescott, R. E. 1969. Ecology of the American dog tick, *Dermacentor variabilis* (Say), in Nova Scotia. Canadian Journal of Zoology 47: 171–181. https://doi.org/10.1139/z69-039

Doran, D. J. 1954a. A catalogue of the protozoa and helminths of North American rodents. I. Protozoa and Acanthocephala. The American Midland Naturalist 52: 118–128.

Doran, D. J. 1954b. A catalogue of the protozoa and helminths of North American rodents. II. Cestoda. The American Midland Naturalist 52: 469–480.

Doran, D. J. 1955a. A catalogue of the protozoa and helminths of North American rodents. III. Nematoda. The American Midland Naturalist 53: 162–175.

Doran, D. J. 1955b. A catalogue of the protozoa and helminths of North American rodents. IV. Trematoda. The American Midland Naturalist 53: 446–454.

Douglas, C. L. 1969. Comparative ecology of pinyon mice and deer mice in Mesa Verde National Park, Colorado. University of Kansas publications, Museum of Natural History 18: 421–504.

Douglass, R. J.; Calisher, C. H.; Wagoner, K. D.; and Mills, J. N. 2007. Sin Nombre virus infection of deer mice in Montana: characteristics of newly infected mice, incidence, and temporal pattern of infection. Journal of

Wildlife Diseases 43: 12–22. https://doi.org/10.7589/0090-3558-43.1.12

Drebot, M. A.; Gavrilovskaya, I.; Mackow, E. R.; Chen, Z.; Lindsay, R.; Sanchez, A. J.; Nichol, S. T.; and Artsob, H. 2001. Genetic and serotypic characterization of Sin Nombre–like viruses in Canadian *Peromyscus maniculatus* mice. Virus Research 75: 75–86. https://doi.org/10.1016/S0168-1702(01)00227-1

Dubey, J. P. 1983. *Sarcocystis peromysci* n. sp. and *S. idahoensis* in deer mouse (*Peromyscus maniculatus*) in Montana. Canadian Journal of Zoology 61: 1180–1182. https://doi.org/10.1139/z83-159

Dunn, L. H.; and Parker, R. R. 1923. Fleas found on wild animals in the Bitterroot Valley, Montana. Public Health Reports 38: 2763–2775.

Durette-Desset, M. C.; Kinsella, J. M.; and Forrester, D. J. 1972. Arguments en faveur de la double origine des nematodes nearctiques du genre *Heligmosomoides* Hall, 1916. Annales de Parasitologie Humaine et Comparée 47: 365–382.

Durette-Desset, M. C. 1974. Nippostrongylinae (Nematoda: Heligmosomidae) nearctiques. Annales de Parasitologie Humaine et Comparee 49: 435–450.

Durette-Desset, M. C.; and Kinsella, J. 2007. A new species of *Heligmosomoides* (Nematoda, Heligmosomidae) parasitic in *Peromyscus maniculatus* (Rodentia, Cricetidae) from Pennsylvania, USA. Acta Parasitologica 52: 342–345.

Duszynski, D. W.; and Anderson, D. R. 1968. *Isospora peromysci* Davis, 1967 in *Peromyscus maniculatus* in northeastern Colorado. The Journal of Parasitology 54: 640.

Dvorak, J.; Otcenasek, M.; and Prokopic, J. 1965. The distribution of adiaspiromycosis. Journal of Hygiene, Epidemiology, Microbiology and Immunology 9: 510–514.

Dyer, W. G.; and Olsen, O. W. 1967a. Biology of *Mastophorus numidica* (Seurat, 1914) Read and Millemann, 1953 (Nematoda: Spiruridae) with a description of the juvenile stages. Proceedings of the Helminthological Society of Washington 34: 98–103.

Dyer, W. G.; and Olsen, O. W. 1967b. Helminths of the deer mouse, *Peromyscus maniculatus*, from northern Colorado. Bulletin of Wildlife Disease Association 3: 35–36.

Dyer, W. G. 1969. A checklist of the protozoa and helminths of the deer mouse, *Peromyscus maniculatus*, in North America. The American Midland Naturalist 81: 258–262. https://doi.org/10.2307/2423671

Eads, R. B.; Menzies, G. C.; and Hightower, B. G. 1956. The ticks of Texas, with notes on their medical significance. Texas Journal of Science 8: 7–24.

Eads, R. B.; and Campos, E. G. 1977. Notes on the flea genus *Megarthroglossus* (Siphonaptera: Hystrighopsyllidae: Anomiopsyllinae), with description of a new species. Journal of Medical Entomology 14: 97–100. https://doi.org/10.1093/jmedent/14.1.97

Eads, R. B.; and Campos, E. G. 1979. Description of a new species of *Catallagia* (Siphonaptera: Hystrichopsyllidae: Neopsyllinae). Journal of Medical Entomology 16: 291–294. https://doi.org/10.1093/jmedent/16.4.291

Eads, R. B.; Campos, E. G.; and Barnes, A. M. 1979. New records for several flea (Siphonaptera) species in the United States, with observations on species parasitizing carnivores in the Rocky Mountain region. Proceedings – Entomological Society of Washington (USA) 81: 38–42.

Easton, E. R.; and Goulding, R. L. 1974. Ectoparasites in two diverse habitats in western Oregon I. *Ixodes* (Acarina: Ixodidae). Journal of Medical Entomology 11: 413–418. https://doi.org/10.1093/jmedent/11.4.413

Easton, E. R. 1975. Ectoparasites in two diverse habitats in western Oregon II. Chiggers (Acari: Trombiculidae). Journal of Medical Entomology 12: 295–298. https://doi.org/10.1093/jmedent/12.3.295

Easton, E. R. 1983. Ectoparasites in two diverse habitats in western Oregon III. Interrelationship of fleas (Siphonaptera) and their hosts. Journal of Medical Entomology 20: 216–219.

Egoscue, H. J. 1966. New and additional host-flea associations and distributional records of fleas from Utah. The Great Basin Naturalist 26: 71–75.

Egoscue, H. J. 1968. A new species of the genus *Stenistomera* (Siphonaptera: Hystrichopsyllidae). Bulletin of the Southern California Academy of Sciences 67: 138–142.

Egoscue, H. J. 1976. Flea exchange between deer mice and some associated small mammals in western Utah. The Great Basin Naturalist 36: 475–480.

Ehrenford, F. A. 1954. Life cycle of *Nematospiroides dubius* Baylis (Nematoda: Heligmosomidae). The Journal of Parasitology 40: 480–481.

Eisen, L.; Dolan, M. C.; Piesman, J.; and Lane, R. S. 2003. Vector competence of *Ixodes pacificus* and *I. spinipalpis* (Acari: Ixodidae), and reservoir competence of the dusky-footed woodrat (*Neotoma fuscipes*) and the deer mouse (*Peromyscus maniculatus*), for *Borrelia bissettii*. Journal of Medical Entomology 40: 311–320. https://doi.org/10.1603/0022-2585-40.3.311

Eisen, L.; Eisen, R. J.; Mun, J.; Salkeld, D. J.; and Lane, R. S. 2009. Transmission cycles of *Borrelia burgdorferi* and *B. bissettii* in relation to habitat type in northwestern California. Journal of Vector Ecology 34: 81–91. https://doi.org/10.1111/j.1948-7134.2009.00010.x

Elliott, L. H.; Ksiazek, T. G.; Rollin, P. E.; Spiropoulou, C. F.; Morzunov, S.; Monroe, M.; Goldsmith, C. S.; Humphrey, C. D.; Zaki, S. R.; Krebs, J. W.; Maupin, G.; Gage, K.; Childs, J. E.; Nichol, S. T.; and Peters, C. J. 1994. Isolation of the causative agent of Hantavirus Pulmonary Syndrome. The American Journal of Tropical Medicine and Hygiene 51: 102–108. https://doi.org/10.4269/ajtmh.1994.51.102

Elzinga, R. J.; and Rees, D. M. 1964. Comparative rates of ectoparasite infestation on deer and harvest mice. Proceedings of the Utah Academy of Sciences, Arts, and Letters 41: 217–220.

Emerson, K. C. 1971. New records of Anoplura from Mexico. Journal of the Kansas Entomological Society 44: 374–377.

Emmons, R. W. 1988. Ecology of Colorado tick fever. Annual Review of Microbiology 42: 49–64. https://doi.org/10.1146/annurev.mi.42.100188.000405

English, M. P.; Kapica, L.; and Maciejewska, J. 1978. On the occurrence of *Microsporum persicolor* in Montreal, Canada. Mycopathologia 64: 35–37. https://doi.org/10.1007/BF00443086

Enright, J. B.; Longhurst, W.; Franti, C. E.; Wright, M. E.; Dutson, V. J.; and Behymer, D. E. 1969. Some observations on domestic sheep and wildlife relationships in Q-fever. Bulletin of the Wildlife Disease Association 5: 276–283. https://doi.org/10.7589/0090-3558-5.3.276

Enright, J. B.; Behymer, D. E.; Franti, C. E.; Dutson, V. J.; Longhurst, W. M.; Wright, M. E.; and Goggin, J. E. 1971a. The behavior of Q fever rickettsiae isolated from wild animals in northern California. Journal of Wildlife Diseases 7: 83–90. https://doi.org/10.7589/0090-3558-7.2.83

Enright, J. B.; Franti, C. E.; Behymer, D. E.; Longhurst, W. M.; Dutson, V. J.; and Wright, M. E. 1971b. *Coxiella burneti* in a wildlife-livestock environment: distribution of Q fever in wild mammals. American Journal of Epidemiology 94: 79–90. https://doi.org/10.1093/oxfordjournals.aje.a121298

Erickson, A. B. 1938a. Parasites of some Minnesota rodents. The Journal of Mammalogy 19: 252–253.

Erickson, A. B. 1938b. Parasites of some Minnesota Cricetidae and Zapodidae and a host catalogue of helminth parasites of native American mice. The American Midland Naturalist 20: 575–589.

Fain, A.; Kok, N. J. J.; Lukoschus, F. S.; and Clulow, F. V. 1971. Notes on the hypopial nymphs phoretic on mammals in Canada with description of a new species (Acarina: Sarcoptiformes). Canadian Journal of Zoology 49: 15–18. https://doi.org/10.1139/z71-004

Fain, A.; and Whitaker Jr., J. O. 1976. *Dermacarus jonesi* sp. n. (acari: Labidophorinae: Glycyphagidae) from *Peromyscus maniculatus* from western North America. The Journal of Parasitology 62: 119–120. https://doi.org/10.2307/3279056

Fain, A.; and Bochkov, A. 2002. On some little known and a new species of Myobiidae (Acari) associated with rodents. Bulletin et Annales de la Société Royale Belge d'Entomologie 138: 95–105.

Farrell, C. E. 1956. Chiggers of the genus *Euschoengastia* (Acarina: Trombiculidae) in North America. Proceedings of the United States National Museum 106: 85–235.

Favorov, M. O.; Kosoy, M. Y.; Tsarev, S. A.; Childs, J. E.; and Margolis, H. S. 2000. Prevalence of antibody to Hepatitis E virus among rodents in the United States. The Journal of Infectious Diseases 181: 449–455. https://doi.org/10.1086/315273

Fay, F. H.; and Rausch, R. L. 1969. Parasitic organisms in the blood of arvicoline rodents in Alaska. The Journal of Parasitology 55: 1258–1265. https://doi.org/10.2307/3277271

Feigley H. P.; and Worley, D. W. 1988. Prevalence of larval *Echinococcus multilocularis* in native Montana small mammals and susceptibility of laboratory rodents to a Montana isolate. Proceedings of the Helminthological Society Washington 55: 105–108.

Ferris, D. H.; Rhoades, H. E.; and Hanson, L. E. 1959. The isolation of *Leptospira hyos* from a new host, *Peromyscus maniculatus*. The Cornell Veterinarian 49: 344–348.

Ferris, G. F. 1916. A catalogue and host list of the Anoplura. Proceedings of the California Academy of Sciences. 4th ser. 6: 120–213.

Ferris, G. F. 1916. Notes on Anoplura and Mallophaga, from mammals with description of four new species and a new variety of Anoplura. Psyche 23: 97–120.

Ferris, G. F. 1921. Contributions towards a monograph of the sucking lice. Part II. Stanford University Publications, University Series, Biological Sciences 2: 52–133.

Ferris, G. F. 1951. The sucking lice. Memoirs of the Pacific Coast Entomological Society 1: 1–320.

Feuer, R.; Boone, J. D.; Netski, D.; Morzunov, S. P.; and Jeor, S. C. S. 1999. Temporal and spatial analysis of Sin Nombre virus quasispecies in naturally infected rodents. Journal of Virology 73: 9544–9554. https://doi.org/10.1128/JVI.73.11.9544-9554.1999

Fish, P. G. 1972. Notes on *Moniliformis clarki* (Ward) (Acanthocephala: Moniliformidae) in West Central Indiana. The Journal of Parasitology 58: 147.

Ford, P. L.; Fagerlund, R. A.; Duszynski, D. W.; and Polechla, P. J. 2004. Fleas and Lice of Mammals in New Mexico. General Technical Rport RMRS-GTR-123. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, Colorado, USA.

Forrester, D. J. 1971. *Heligmosomoides polygyrus* (= *Nematospiroides dubius*) from wild rodents of northern California: natural infections, host specificity, and strain characteristics. The Journal of Parasitology 57: 498–503. https://doi.org/10.2307/3277902

Forrester, D. J.; and Neilson, J. T. McL. 1973. Comparative infectivity of *Heligmosomoides polygyrus* (= *Nematospiroides*) in three species of *Peromyscus*. The Journal of Parasitology 59: 251–255.

Fox, I. 1940. Fleas of Eastern United States. Ames: Iowa State College Press. 191 pp.

Frandsen, J. C.; and Grundmann, A. W. 1960. *Brachylaime microti* from the deer mouse in Utah. The Journal of Parasitology 46: 314.

Frandsen, J. C.; and Grundmann, A. W. 1961a. *Trichuris stansburyi* and *Gongylonema mysciphilia*, two new species of nematodes from the deer mouse in Utah. Proceedings of the Helminthological Society of Washington 28: 91–94.

Frandsen, J. C.; and Grundmann, A. W. 1961b. Endoparasitism in isolated populations of rodents of the Lake Bonneville Basin, Utah. The Journal of Parasitology 47: 391–396.

Freeman, R. S. 1956. Life history studies on *Taenia mustelae* Gmelin, 1700 and the taxonomy of certain taenoid cestodes from Mustelidae. Canadian Journal of Zoology 34: 219–212.

Freeman, R. S. 1958. On the epizootiology of *Capillaria hepatica* (Bancroft, 1893) in Algonquin Park, Ontario. The Journal of Parasitology 44: 33.

Freeman, R. S. 1960. Another hymenolepid with great morphological variation: *Hymenolepis bennetti* n. sp. (Cestoda) from *Napaeozapus insignis algonquinensis* Prince. Canadian Journal of Zoology 38: 737–743.

Freeman, R. S.; and Wright, K. A. 1960. Factors concerned with the epizootiology of *Capillaria hepatica* (Bancroft, 1893) (Nematoda) in a population of *Peromyscus maniculatus* in Algonquin Park, Canada. The Journal of Parasitology 46: 373–382.

Frenkel, J. K.; and Lunde, M. N. 1966. Effects of corticosteroids on antibody and immunity in "Besnoitia" infection of hamsters. The Journal of Infectious Diseases 116: 414–424.

Fuller, C. A. 1996. Population dynamics of two species of *Eimeria* (Apicomplexa: Eimeriidae) in deer mice (*Peromyscus maniculatus*): biotic and abiotic factors. The Journal of Parasitology 82: 220–225. https://doi.org/10.2307/3284150

Furman, D. P. 1955. Revision of the genus *Eubrachylaelaps* (Acarina: Laelaptidae) with the descriptions of two new species. Annals of the Entomological Society of America 18: 51–59.

Garvie, M. B.; Mckiel, J. A.; Sonenshine, D. E.; and Campbell, A. 1978. Seasonal dynamics of American dog tick, *Dermacentor variabilis* (Say), populations in southwestern Nova Scotia. Canadian Journal of Zoology 56: 28–39. https://doi.org/10.1139/z78-004

Gastfriend, A. 1955. New host records for the immature stages of the tick *Dermacentor parumapertus*. The Journal of Parasitology 41: 63–65.

Geary, J. M. 1959. The fleas of New York. Cornell University Agricultural Experimental Station Memoir 355, Ithaca, New York. 104 pp.

Giesen, K. M. T.; Lukoschus, F. S.; Whitaker Jr., J. O.; and Gettinger, D. 1983. Four new species of itch mites (Acari: Psorergatidae: Prostigmata) from small mammals in North America. Journal of Medical Entomology 20: 164–173. https://doi.org/10.1093/jmedent/20.2.164

Glass, G. E.; Yates, T. L.; Fine, J. B.; Shields, T. M.; Kendall, J. B.; Hope, A. G.; Parmenter, C. A.; Peters, C. J.; Ksiazek, T. G.; Li, C. S.; Patz, J. A.; and Mills, J. N. 2002. Satellite imagery characterizes local animal reservoir populations of Sin Nombre virus in the southwestern United States. Proceedings of the National Academy of Sciences 99: 16817–16822. https://doi.org/10.1073/pnas.252617999

Glicken, A.; and Schwab, R. G. 1980. Modes of ectoparasite reinfestations of deer mice (*Peromyscus maniculatus*). Journal of Wildlife Diseases 16: 577–586. https://doi.org/10.7589/0090-3558-16.4.577

Goff, M. L.; and Loomis, R. B. 1973. Two new species of *Odontacarus* Ewing (Acarina: Trombiculidae) from California and Baja California, Mexico. Journal of Medical Entomology 10: 333–336. https://doi.org/10.1093/jmedent/10.4.333

Goff, M. L.; and Loomis, R. B. 1974. New species of *Comatacarus* (Acarina: Trombiculidae) from central United States. Bulletin of the Southern California Academy of Sciences 73: 170–171.

Goff, M. L.; Loomis, R. B.; and Brennan, J. M. 1972. *Odontacarus dentatus* (Ewing, 1925) is the senior synonym of *O. galli* (Ewing, 1946) (Acarina: Trombiculidae). Journal of Medical Entomology 9: 479–481. https://doi.org/10.1093/jmedent/9.5.479

Goodrich, I.; McKee, C.; and Kosoy, M. 2020. Longitudinal study of bacterial infectious agents in a community of small mammals in New Mexico. Vector-Borne and Zoonotic Diseases 20: 496–508. https://doi.org/10.1089/vbz.2019.2550

Gould, D. J. 1956. The larval trombiculid mites of California (Acarina, Trombiculidae). University of California Publications in Entomology 11: 1–116.

Graham, T. B.; and Chomel, B. B. 1997. Population dynamics of the deer mouse (*Peromyscus maniculatus*) and Sin Nombre virus, California Channel Islands. Emerging Infectious Diseases 3: 367–370.

Grant, D. R.; and Woo, P. T. K. 1979. Comparative studies of *Giardia* spp. in small mammals in southern Ontario. III. Duration and cyst production in natural and experimental infections. Canadian Journal of Zoology 57: 307–313. https://doi.org/10.1139/z79-034

Greenberg, B. 1951. A new subgenus of *Acomatacarus* from Kansas. (Acarina, Trombiculidae). The Journal of Parasitology 37: 525–527.

Greenberg, B. 1952. A review of the New World *Acomatacarus* (Acarina, Trombiculidae). Annals of the Entomological Society of America 45: 473–491.

Grundmann, A. W. 1956. A new tapeworm, *Mesocestoides carnivoricolus*, from carnivores of the Great Salt Lake Desert region of Utah. Proceedings of the Helminthological Society of Washington 23: 26–28.

Grundmann, A. W. 1957a. Nematode parasites of mammals of the Great Salt Lake Desert of Utah. The Journal of Parasitology 43: 105–112.

Grundmann, A. W. 1957b. Contribution of parasitism studies to the elucidation of food cycles of some local mammals. Proceedings of the Utah Academy of Sciences, Arts, and Letters 34: 61–63.

Grundmann, A. W., 1958. Cestodes of mammals from the Great Salt Lake desert region of Utah. The Journal of Parasitology 44: 425–429.

Grundmann, A. W. 1959. A study of parasitism of the deer mouse, *Peromyscus maniculatus*, in the Bonneville Basin of Utah and its role in parasite distribution. The Journal of Parasitology 45 (supplement): 34.

Grundmann, A. W.; and Frandsen, J. C. 1959. A study of the parasitism of the deer mouse, *Peromyscus maniculatus*, in the Bonneville Basin of Utah and its role in parasite distribution. The Journal of Parasitology 45: 34–35.

Grundmann, A. W.; and Frandsen, J. C. 1960. Definitive host relationships of the helminth parasites of the deer mouse, *Peromyscus maniculatus*, in the Bonneville Basin of Utah. The Journal of Parasitology 46: 673–677.

Grundmann, A. W.; Frandsen, J. C.; and Warnock, R. C. 1961. A study of geographical isolation of parasitic nematodes on Stansbury Island, Great Salt Lake. Proceedings of the Utah Academy of Sciences, Arts, and Letters 38: 74–77.

Grundmann, A. W.; and Warnock, R. C. 1964. A population study of eight parasitic nematode species from mammals of the Bonneville Basin of Utah. Proceedings of the Utah Academy of Sciences, Arts, and Letters 41: 228–233.

Grundmann, A. W.; Warnock, R. C.; and Wassom, D. L. 1976. Some mechanisms of natural regulation of parasitic helminth populations. The American Midland Naturalist 95: 347–360.

Guilday, J. E. 1962. The Pleistocene local fauna of the natural chimneys, Augusta County, Virginia. Annals of the Carnegie Museum 36: 87–122. Cited by Hibbard (1968).

Guilday, J. E.; Martin, P. S.; and McCrady, A. D. 1964. New Paris no. 4: a late Pleistocene cave deposit in Bedford County, Pennsylvania. Bulletin of the National Speleological Society 26: 122–194. Cited by Hibbard (1968).

Haas, G. E.; Martin, R. P.; Swickard, M.; and Miller, B. E. 1973. Siphonaptera-mammal relationships in northcentral New Mexico. Journal of Medical Entomology 10: 281–289. https://doi.org/10.1093/jmedent/10.3.281

Haas, G. E.; Johnson, L. and Wilson, N. 1980. Siphonaptera from mammals in Alaska. Supplement II. Southeastern Alaska. Journal of the Entomological Society of British Columbia 77: 43–46.

Haley, A. J. 1961. Biology of the rat nematode *Nippostrongylus brasiliensis* (Travassos, 1914). I. Systematics, host and geographical distribution. The Journal of Parasitology 47: 727–732.

Hall, M. C. 1912. The parasite fauna of Colorado. Colorado College Publication, Science Series 12: 329–383.

Hall, M. C. 1916. Nematode parasites of mammals of the orders Rodentia, Lagomorpha and Hyracoidea. Proceedings of the United States National Museum 50: 1–258.

Hall, J. E.; and Sonnenberg, B. 1955. Some helminth parasites of rodents from localities in Maryland and Kentucky. The Journal of Parasitology 41: 640–641. (No parasites from *P. maniculatus* reported.)

Hamilton, W. J. 1941. The food of small forest mammals in eastern United States. The Journal of Mammalogy 21: 250–263.

Hansen, C. G. 1964. Ectoparasites of mammals from Oregon. Great Basin Naturalist 24: 75–81.

Hansen, M. F. 1950. A new dilepidid tapeworm and notes on other tapeworms of rodents. The American Midland Naturalist 43: 471–479.

Hardy, J. L.; Scrivani, R. P.; Lyness, R. N.; Nelson, R. L.; and Boberts, D. R. 1970. Ecologic studies on Buttonwillow virus in Kern County, California, 1961–1968. The American Journal of Tropical Medicine and Hygiene 19: 552–563. https://doi.org/10.4269/ajtmh.1970.19.552

Hardy, J. L.; Reeves, W. C.; Scrivani, R. P.; and Roberts, D. R. 1974. Wild mammals as hosts of group A and group B arboviruses in Kern County, California. The American Journal of Tropical Medicine and Hygiene 23: 1165–1177. https://doi.org/10.4269/ajtmh.1974.23.1165

Harkema, R. 1936. The parasites of some North Carolina rodents. Ecological Monographs 6: 151–232.

Harvey, R. V. 1907. British Columbia fleas. Bulletin of the Entomological Society of British Columbia 7: 1–4.

Healey M. C.; and Grundmann, A. W. 1974. The influence of intermediate hosts on the infection pattern of *Protospirura numidica criceticola* Quentin, Karimi, and Rodriguez De Almeida, 1968 (Nematoda: Spiruridae) in the Bonneville Basin, Utah. Proceedings of the Helminthological Society of Washington 41: 59–63.

Hedeen, R. A. 1953. Ectoparasites occurring on mammals in the vicinity of Ft. Hood, Texas. Texas Journal of Science 5: 125–129.

Herman, T.B. 1981. *Capillaria hepatica* (Nematoda) in insular populations of the deer mouse *Peromyscus maniculatus*: cannibalism or competition for carcasses? Canadian Journal of Zoology 59: 776–784.

Herrin, C. S. 1970. A systematic revision of the genus *Hirstionyssus* (Agari: Mesostigmata) of the Nearctic Region. Journal of Medical Entomology 7: 391–437. https://doi.org/10.1093/jmedent/7.4.391

Hibbard, C. W. 1968. Palaeontology. Pp. 6–26 in: Biology of *Peromyscus* (Rodentia), King, J. A. (ed.), Special Publication No 2. The American Society of Mammalogists 1968. 593 pp.

Holdenried, R.; Evans, F. C.; and Longanecker, D. S. 1951. Host-parasite-disease relationships in a mammalian community in the central coast range of California. Ecological Monographs 21: 1–18.

Holdenried, R.; and Morlan, H. B. 1955. Plague-infected fleas from northern New Mexico wild rodents. The Journal of Infectious Diseases 96: 133–137.

Holdenried, R.; and Morlan, H. B. 1956. A field study of wild mammals and fleas of Santa Fe County, New Mexico. The American Midland Naturalist 55: 369–381.

Holdenried, R.; and Quan, S. F. 1956. Susceptibility of New Mexico rodents to experimental plague. Public Health Reports 71: 979–984.

Holland, G. P. 1941. Further records of Siphonaptera for British Columbia. Journal of the Entomological Society of British Columbia 37: 10–14.

Holland, G. P. 1944. Notes on some northern Canadian Siphonaptera, with the description of a new species. The Canadian Entomologist 76: 212–246.

Holland, G. P. 1949. The Siphonaptera of Canada. Publication 817, Technical Bulletin 70. Dominion of Canada, Department of Agriculture, Kamloops, British Columbia.

Holland, G. P.; and Benton, A. H. 1968. Siphonaptera from Pennsylvania mammals. The American Midland Naturalist 80: 252–261. https://doi.org/10.2307/2423612

Holland, G. P. 1979. Three new species of fleas from Canada (Siphonaptera). The Canadian Entomologist 111: 713–719. https://doi.org/10.4039/Ent111713-6

Holloway, H. L. 1963. Observations on the morphology and life cycle of a cestode possessing small hooks with bifid guards. The Journal of Parasitology 49: 23–24.

Holmes, J. C.; Mahrt, J. L.; and Samuel, W. M. 1971. The occurrence of *Echinococcus multilocularis* Leuckart, 1863 in Alberta. Canadian Journal of Zoology 49: 575–576.

Howell, J. F.; and Elzinga, R. J. 1962. A new *Radfordia* (Acarina: Myobiidae) from the kangaroo rat and a key to the known species. Annals of the Entomological Society of America 55: 547–555.

Howell, L.; Jelden, K.; Rácz, S. E.; Gardner, S. L.; and Gettinger, D. 2016. Arthropods infesting small mammals (Insectivora and Rodentia) near Cedar Point Biological Station in southwestern Nebraska. Insecta Mundi 0478: 1–16. https://digitalcommons.unl.edu/insectamundi/954/

Hubálek, Z. 2000. Keratinophilic fungi associated with free-living mammals and birds. Revista Iberoamericana de Micología 17: 93–103.

Hubálek, Z.; and Rudolf, I. 2011. Vertebrates as hosts and reservoirs of zoonotic microbial agents. Pp. 83–128 in: Microbial Zoonoses and Sapronoses, Hubálek, Z.; and Rudolf, I. (eds.) Springer Netherlands, Dordrecht.

Hubbard, C. A. 1947. The Fleas of Western North America. Iowa State College Press. 533 pp.

Hubbard, C. A. 1949. Fleas of the state of Nevada. Bulletin of the Southern California Academy of Sciences 48: 115–128.

Hubbard, D. H. 1941. Some mammalian ectoparasites. The Journal of Mammalogy 22: 202–203.

Hubbert, W. T.; Goldenberg, M. I.; Kartman, L.; and Prince, F. M. 1966. Public health potential of sylvatic plague. Journal of the American Veterinary Medical Association 149: 1651–1654.

Hudson, B. W.; Quan, S. F.; and Goldenberg, M. I. 1964. Serum antibody responses in a population of *Microtus californicus* and associated rodent species during and after *Pasteurella pestis* epizootics in the San Francisco Bay area. Zoonoses Research 3: 15–29.

Hudson, B. W.; Goldenberg, M. I.; and Quan, T. J. 1972. Serologic and bacteriologic studies on the distribution of plague infection in a wild rodent plague pocket in the San Francisco Bay area of California. Journal of Wildlife Diseases 8: 278–286. https://doi.org/10.7589/0090-3558-8.3.278

Hudson, B. W.; and Quan, T. J. 1975. Serologic observations during an outbreak of rat borne plague in the San Francisco Bay area of California. Journal of Wildlife Diseases 11: 431–436. https://doi.org/10.7589/0090-3558-11.3.431

Hume, G.; Hawkinson, A.; Aboellail, T.; and Schountz, T. 2015. Experimental Modoc virus infection of deer mice (*Peromyscus maniculatus*). New Horizons in Translational Medicine 2: 128–129. https://doi.org/10.1016/j.nhtm.2015.07.048

Hunninen, A. V. 1935. Infection of abnormal hosts with the mouse strain of *Hymenolepis fraterna*. The Journal of Parasitology 21: 312.

Hunter, W. D.; and Bishopp, F. C. 1911. The Rocky Mountain spotted fever tick, with special reference to the problem of its control in the Bitterroot Valley in Montana. USDA Bureau of Entomology, Bulletion No. 105. 47 pp.

Hunter, D. M.; Sadleir, R. M. F. S.; and Webster, J. M. 1972. Studies on the ecology of cuterebrid parasitism in deermice. Canadian Journal of Zoology 50: 25–29. https://doi.org/10.1139/z72-005

Hwang, Y. T.; Gardner, S. L.; and Millar, J. S. *Hymenolepis horrida* (Cestoda: Hymenolepididae) and *Catenotaenia peromysci* (Cestoda: Anoplocephalidae) in voles from the Canadian Rockies. Comparative Parasitology 74: 160–163.

Jacobsen, B. R. 1968. The helminthofauna of small rodents in southern Manitoba. Master's thesis, University of Manitoba, Winnipeg.

Jameson, E. W. 1952. Food of deer mice, *Peromyscus maniculatus* and *P. boylei*, in the northern Sierra Nevada, California. The Journal of Mammalogy 33: 50–60. https://doi.org/10.2307/1375640

Jameson Jr., E. W. 1950. *Hirstionyssus obsoletus*, a new mesostigmatic mite from small mammals of the western United States. Proceedings of the Biological Society of Washington 63: 31–34.

Jameson Jr., E. W.; and Brennan, J. M. 1957. An environmental analysis of some ectoparasites of small forest mammals in the Sierra Nevada, California. Ecological Monographs 27: 45–54.

Jellison, W. I. 1971. Adiaspiromycosis. Pp. 321–322 in: Infectious diseases of wild mammals, Davis, J. W.; Karstad, L. H.; and Trainer, D. O. (eds.). Iowa State University Press, Ames.

Jellison. W. L. 1939. *Opisodasys* Jordan, 1933, a genus of Siphonaptera. The Journal of Parasitology 25: 413–420.

Jellison, W. L.; and Good, N. E. 1942. Index to the literature of Siphonaptera of North America. National Institute of Health Bulletin 178: 1–193.

Johnson, D. W. 1966. Ticks of Dugway Proving Ground and vicinity and their host associations. Proceedings of the Utah Academy of Sciences, Arts, and Letters 43: 49–66.

Johnson, H. N. 1970. Keynote address: the ecological approach to the study of zoonotic diseases. Journal of Wildlife Diseases 64: 194–204.

Jones, A.; and Pybus, M. J. 2001. Taeniasis and echinococcus. Pp. 150–192 in: Parasitic Diseases of Wild Mammals, Samuel, W. M.; Pybus, M. J.; and Kocan, A. A. (eds.). Iowa State University Press, Ames.

Jordan, R. 1928. Siphonaptera collected during a visit to the eastern United States of North America in 1927. Novitates Zoologicae 34: 178–188.

Jordan, K. 1929. Notes on North American fleas. Novitates Zoologicae 35: 28–39.

Jordan, R. 1932. Siphonaptera collected by Mr. Harry S. Swarth at Atlin in British Columbia. Novitates Zoologicae 38: 253–255.

Judd, W. W. 1955. A collection of fleas from the vicinity of Fort Simpson, Northwest Territories, Canada. The Journal of Parasitology 41: 441–442.

Kapoor, A.; Simmonds, P.; Scheel, T. K. H.; Hjelle, B.; Cullen, J. M.; Burbelo, P. D.; Chauhan, L. V.; Duraisamy, R.; Leon, M. S.; Jain, K.; Vandegrift, K. J.; Calisher, C. H.; Rice, C. M.; and Lipkin, W. I. 2013. Identification of rodent homologs of hepatitis C virus and pegiviruses. mBio 4. https://doi.org/10.1128/mBio.00216-13

Karabatsos, N. (Ed.) 1985. International catalogue of arboviruses, including certain other viruses of vertebrates. 3rd ed. American Society of Tropical Medicine and Hygiene for the Subcommittee on Information Exchange of the American Committee on Arthropod-borne Viruses, San Antonio, Texas.

Kardos, E. H. 1954. Biological and systematic studies on the subgenus *Neotrombicula* (genus *Trombicula*) in the central United States (Acarina: Trombiculidae). The University of Kansas Science Bulletin 36 (part 1): 69–123.

Kartman, L.; Quan, S. F.; and Stark, H. E. 1962. Ecological studies of wild rodent plague in the San Francisco Bay area of California. VII. Effects of plague in nature on *Microtus californicus* and other wild rodents. Zoonoses Research 1: 99–119.

Keegan, H. L. 1951. The mites of the subfamily Haemogamasinae (Acari: Laelaptidae). Proceedings of the United States National Museum 101: 203–268.

Keegan, H. L. 1953. Collections of parasitic mites from Utah. The Great Basin Naturalist 13: 35–42.

Keegan, H. L. 1956. *Ischyropoda furmani* n. sp., a new ectoparasitic mite from Utah. The Journal of Parasitology 42: 311–315.

Kegley, L. M.; Baldwin, J.; Brown, B. W.; and Berntzen, A. K. 1970. *Mesocestoides corti*: Environmental cation concentration in calcareous corpuscles. Experimental Parasitology 27: 88–94.

Kellogg, V. L.; and Ferris, G. F. 1915. The Anoplura and Mallogphaga of North American Mammals. Leland Stanford Junior University Publications. University Series no. 20. Stanford University, California. 74 pp.

Kim, R. C. 1965. A review of the *Hoplopleura hesperomydis* complex (Anoplura: Hoplopleuridae). The Journal of Parasitology 51: 871–887.

Kim, K. C.; Brown Jr., B. W.; and Cook, E. F. 1966. A quantitative taxonomic study of the *Hoplopleura hesperomydis* complex (Anoplura, Hoplopleuridae), with notes on *A. posteriori* taxonomic characters. Systematic Biology 15: 24–45. https://doi.org/10.2307/sysbio/15.1.24

King, J. A. 1968. Biology of *Peromyscus* (Rodentia). Special Publication no. 2. The American Society of Mammalogists, Stillwater, Oklahoma. 616 pp. https://doi.org/10.5962/bhl.title.39510

Kinsella, J. M.; and Pattie, D. L. 1967. Ectoparasites of small mammals of the alpine Beartooth Plateau, Wyoming. Canadian Journal of Zoology 45: 233–235. https://doi.org/10.1139/z67-031

Kinsella, J. M. 1968. *Monopsyllus thambus* (Jordan, 1929) from the alpine in Colorado. The Journal of Parasitology 54: 258.

Kinsella, J. M. 1987. Studies on the life cycle and host specificity of *Parastrongylus schmidti* (Nematoda: Angiostrongylidae) Proceedings of the Helminthological Society of Washington 54: 245–248.

Kinsella J. M. 1971. Growth, development, and intraspecific variation of *Quinqueserialis quinqueserialis* (Trematoda: Notocootylidae) in rodent hosts. The Journal of Parasitology 57: 62–70.

Koehler, D. K.; and Anderson, S. H. 1991. Habitat use and food selection of small mammals near a sagebrush/crested wheatgrass interface in southeastern Idaho. The Great Basin Naturalist 51: 249–255.

Kofoid, C. A.; and Christiansen, E. B. 1915a. On the life-history of *Giardia*. Proceedings of the National Academy of Sciences of the United States of America 1: 547–552.

Kofoid, C. A.; and Christiansen, E. B. 1915b. On binary and multiple fission in *Giardia muris* (Grassi). University of California Publications in Zoology 16: 30–54.

Kohls, G. M.; and Clifford, C. M. 1963. *Ornithodoros sparnus* sp. n., a parasite of wood rats, *Neotoma* spp., and deer mice, *Peromyscus* spp., in Utah and Arizona (Acarina: Argasidae). The Journal of Parasitology 49: 857–861. https://doi.org/10.2307/3275937

Kohls, G. M.; Sonenshine, D. E.; and Clifford, C. M. 1965. The systematics of the Subfamily Ornithodorinae (Acarina: Argasidae). II. Identification of the larvae of the western hemisphere and descriptions of three new species. Annals of the Entomological Society of America 58: 331–364. https://doi.org/10.1093/aesa/58.3.331

Kok, N. J. J.; Lukoschus, F. S.; and Clulowi, F. V. 1971. Three new itch mites from Canadian small mammals (Acarina: Psorergatidae). Canadian Journal of Zoology 49: 1239–1248. https://doi.org/10.1139/z71-187

Kosoy, M. Y.; Regnery, R. L.; Tzianabos, T.; Marston, E. L.; Jones, D. C.; Green, D.; Maupin, G. O.; Olson, J. G.; and Childs, J. E. 1997. Distribution, diversity, and host specificity of *Bartonella* in rodents from the southeastern United States. The American Journal of Tropical Medicine and Hygiene 57: 578–588. https://doi.org/10.4269/ajtmh.1997.57.578

Kritsky, D. C.; and Leiby, P. D. 1975. Comparison of yearly prevalences of *Echinococcus multilocularis* Leuckart 1863 in *Peromyscus maniculatus* and *Microtus pennsylvanicus* in North Dakota. The Journal of Parasitology 61: 1112–1113.

Kruidenier, F. J.; and Peebles, C. R. 1957. Nematodes in Grand Canyon rodents: *Syphacia*, *Gongylonema*, and *Rictularia*. The Journal of Parasitology 43: 29.

Kruidenier, F. J.; and Peebles, C. R. 1958. *Gongylonema* of rodents: *G. neoplasticum* (redefinition); *G. dipodomysis* n. sp.; and *G. peromysci* n. sp. Transactions of the American Microscopical Society 77: 307–315.

Kruidenier, F. J.; Mehra, K. N.; and Harkema, R. 1961. Comparative studies of *Syphacia peromysci* and *S. samorodini* (Nematoda: Oxyuridae). The Journal of Parasitology 47: 47–51.

Kuenzi, A. J.; Douglass, R. J.; Bond, C. W.; Calisher, C. H.; and Mills, J. N. 2005. Long-term dynamics of Sin Nombre viral RNA and antibody in deer mice in Montana. Journal of Wildlife Diseases 41: 473–481. https://doi.org/10.7589/0090-3558-41.3.473

Lane, R. S.; Emmons, R. W.; Dondero, D. V.; and Nelson, B. C. 1981. Ecology of tick-borne agents in California. I. Spotted fever group rickettsiae. The American Journal of Tropical Medicine and Hygiene 30: 239–252.

Lane, R. S.; Emmons, R. W.; Devlin, V.; Dondero, D. V.; and Nelson, B. C. 1982. Survey for evidence of Colorado tick fever virus outside of the known endemic area in California. The American Journal of Tropical Medicine and Hygiene 31: 837–843. https://doi.org/10.4269/ajtmh.1982.31.837

Lankester, M. W.; and Anderson, R. C. 1966. Small mammals as paratenic hosts of lungworms. Canadian Journal of Zoology 44: 341–342.

Larson, S. R.; Lee, X.; and Paskewitz, S. M. 2018. Prevalence of tick-borne pathogens in two species of *Peromyscus* mice common in northern Wisconsin. Journal of Medical Entomology 55: 1002–1010. https://doi.org/10.1093/jme/tjy027

Lawrence, W. H.; Hays, K. L.; and Graham, S. A. 1965. Arthropodous ectoparasites from some northern Michigan mammals. Occasional Papers of the Museum of Zoology, University of Michigan 639: 1–7.

Layne, J. N. 1958. Records of fleas (Siphonaptera) from Illinois mammals. Natural History Miscellanea, no. 162. Chicago Academy of Sciences, Chicago.

Layne, J. N. 1968. Host and ecological relationships of the parasitic helminth *Capillaria hepatica* in Florida mammals. Zoologica: New York Zoological Society 53: 107–123.

Lee, C. 1969. Larval *Echinococcus multilocularis* Leuckart, 1863 in the southern Interlake area of Manitoba. Canadian Journal of Zoology 47: 733–734.

Leiby, P. D. 1961. Intestinal helminths of some Colorado mammals. The Journal of Parasitology 47: 311.

Leiby, P. D. 1962. Helminth parasites recovered from some rodents in southwestern Idaho. The American Midland Naturalist 67: 250.

Leiby, P. D. 1965. Cestode in North Dakota: *Echinococcus* in field mice. Science 150: 763.

Leiby, P. D.; Carney, W. P.; and Woods, C. E. 1970. Studies on sylvatic echinococcosis. III. Host occurrence and geographic distribution of *Echinococcus multilocularis* in the north central United States. The Journal of Parasitology 56: 1141–1150.

Leiby, P. D.; and Kritsky, D. C. 1972. *Echinococcus multilocularis*: a possible domestic life cycle in central North America and its public health implications. The Journal of Parasitology 58: 1213–1215.

Leiby, P. D.; and Kritsky, D. C. 1974. Studies on sylvatic echinococcosis. IV. Ecology of *Echinococcus multilocularis* in the intermediate host, *Peromyscus maniculatus*, in North Dakota, 1965–1972. The American Journal of Tropical Medicine and Hygiene 23: 667–675.

Leiby, P. D.; Lubinsky, G.; and Galaugher, W. 1969. Studies on sylvatic echinococcosis. II. The occurrence of *Echinococcus multilocularis* Leuck., 1863 in Manitoba. Canadian Journal of Zoology 47: 135–139.

Leiby, P. D.; and Nickel, M. P. 1968. Studies on sylvatic echinococcosis. I. Ground beetle transmission of *Echinococcus multilocularis* Leuckart, 1863, to deer mice, *Peromyscus maniculatus* (Wagner). The Journal of Parasitology 54: 536–737.

Leiby, D. A.; Schad, G. A.; Duffy, C. H.; and Murrell, K. D. 1988. *Trichinella spiralis* in an agricultural ecosystem. III. Epidemiological investigations of *Trichinella spiralis* in resident wild and feral animals. Journal of Wildlife Diseases 24: 606–609. https://doi.org/10.7589/0090-3558-24.4.606

Levine, N. D.; Ivens, V.; and Kruidenier, F. J. 1955. Two new species of *Klossia* (Sporozoa: Adeleidae) from a deer mouse and a bat. The Journal of Parasitology 41: 623–629.

Levine, N. D. and Ivens, V. 1963. *Eimeria siniffi* sp. n. and *E. arizonensis* (Protozoa : Eimeriidae) from deermice in British Columbia. The Journal of Parasitology 49: 660–661. https://doi.org/10.2307/3275782

Lewis, R. E.; and Maser, C. 1978. *Phalacropsylla oregonensis* sp. n., with a key to the species of *Phalacropsylla* Rothschild 1915 (Siphonaptera: Hystrichopsyllidae). The Journal of Parasitology 64: 147–150. https://doi.org/10.2307/3279629

Lichtenfels, J. R. 1966. A survey of the intestinal nematodes of some Maryland rodents with a host-parasite catalog. Master's thesis, University of Maryland. 76 pp.

Lichtenfels, J. R. 1970. Two new species of *Pterygodermatites* (Paucipectines) Quentin, 1969 (Nematoda: Rictulariidae) with a key to the species from North American rodents. Proceedings of the Helminthological Society of Washington 37: 94–101.

Lipovsky, L. J. 1951a. A new genus of Walchiinae (Acarina: Trombiculidae). Journal of the Kansas Entomological Society 24: 95–102.

Lipovsky, L. J. 1951b. A washing method of ectoparasite recovery with particular reference to chiggers (Acarina: Trombiculidae). Journal of the Kansas Entomological Society 24: 151–156.

Lipovsky, L. J.; and Loomis, R. B. 1954. A new chigger mite genus *Euschoengastia* from the central United States. The Journal of Parasitology 40: 107–109.

Llewellyn, J. R. 1978. Differential parasitism of *Peromyscus maniculatus* and *Peromyscus truei* by *Cuterebra* larvae. Great Basin Naturalist 38: 51–54.

Locker, B. 1955. The identification of *Taenia tenuicollis* Rudolphi, 1819 in North America. The Journal of Parasitology 41: 51–56.

Logiudice, K. 2001. Latrine foraging strategies of two small mammals: implications for the transmission of *Baylisascaris procyonis*. The American Midland Naturalist 146: 369–378.

Loomis, R. B. 1954. A new subgenus and six new species of chigger mites (genus *Trombicula*) from the central United States. The University of Kansas Science Bulletin 36: 919–941.

Loomis, R. B.; and Lipovsky, L. J. 1954. Two new chigger mites (genus *Trombicula*) from the central United States. The Journal of Parasitology 27: 47–53.

Loomis, R. B. 1955. *Trombicula gurneyi* Ewing and two related chigger mites (Acarina: Trombiculidae). The University of Kansas Science Bulletin 37: 251–267.

Loomis R. B. 1956. The chigger mites of Kansas (Acarina: Trombiculidae). The University of Kansas Science Bulletin 37: 1195–1443.

Loomis, R. B.; and Bunnell, D. M. 1962. A new species of chigger, genus *Euschoengastia* (Acarina: Trombiculidae), with notes on other species of chiggers from the Santa Ana Mountains, California. Bulletin of the Southern California Academy of Sciences 61: 177–181.

Loomis, R. B. 1971. The genus *Euschoengastoides* (Acarina: Trombiculidae) from North America. The Journal of Parasitology 57: 689–707. https://doi.org/10.2307/3277788

Loomis, R. B.; and Goff, M. L. 1973. A new genus and two new species of North American leeuwenhoekine chiggers (Acarina: Trombiculidae). Journal of Medical Entomology 10: 113–117. https://doi.org/10.1093/jmedent/10.2.113

Lubinsky, G. 1956. On the probable presence of parasitic liver cirrhosis in Canada. Canadian Journal of Comparative Medicine and Veterinary Science 20: 457–465.

Lubinsky, G. 1957. List of helminths from Alberta rodents. Canadian Journal of Zoology 35: 623–627.

Lubinsky G.; Jacobsen, B. R.; and Baron, R. W. 1971. Wildlife foci of *Capillaria hepatica* infections in Manitoba. Canadian Journal of Zoology 49: 1201–1202.

Lundgren, D. L.; and Thorpe, B. D. 1966. Infectious diseases in wild animals in Utah. VII. Experimental infection of rodents with *Rickettsia rickettsii*. The American Journal of Tropical Medicine and Hygiene 15: 799–806. https://doi.org/10.4269/ajtmh.1966.15.799

Madhav, N. K.; Wagoner, K. D.; Douglass, R. J.; and Mills, J. N. 2007. Delayed density-dependent prevalence of Sin Nombre virus antibody in Montana deer mice (*Peromyscus maniculatus*) and implications for human disease risk. Vector-Borne and Zoonotic Diseases 7: 353–364. https://doi.org/10.1089/vbz.2006.0605

Madison, D. M. 1977. Movements and habitat use among interacting *Peromyscus leucopus* as revealed by radiotelemetry. The Canadian Field-Naturalist 91: 273–281.

Main, A. J. 1970. Distribution, seasonal abundance and host preference of fleas in New England. Proceedings of the Entomological Society of Washington 72: 73–89.

Malek, E. A. 1977. Geographical distribution, hosts, and biology of *Schistosomatium douthitti* (Cort, 1914) Price, 1931. Canadian Journal of Zoology 55: 661–671.

Maurer Jr., F. W.; and Skaley, J. E. 1968. Cuterebrid infestation of *Microtus* in eastern North Dakota, Pennsylvania, and New York. Journal of Mammalogy 49: 773–774.

Mathies Jr., A. W. 1958. Certain aspects of the host-parasite relationship of *Aspiculuris tetraptera*, a mouse pinworm. Dissertation Abstracts 18: 1544-1545.

Mayberry, L. F.; Bristol, J. R.; and Duszynski, D. W. 1980. *Isospora californica* (Protozoa: Eimeriidae) in *Peromyscus maniculatus* (Cricetidae) from White Sands National Monument, New Mexico. Southwestern Naturalist 25: 126.

Mayberry, L. F.; Canaris, A. G.; Bristol, J. R.; and Gardner, S. L. 2000. Bibliography of Parasites and Vertebrate Hosts in Arizona, New Mexico and Texas (1893–1984). University of Nebraska Harold W. Manter Laboratory of Parasitology. https://digitalcommons.unl.edu/parasitologyfacpubs/2/

MacDonald, S. O.; Waltari, E.; Nofchissey, R. A.; Sawyer, Y. E.; Ebel, G. D.; and Cook, J. A. 2009. First records of deermice (*Peromyscus maniculatus*) in the Copper River Basin, south-central Alaska. Northwestern Naturalist 90: 243–247.

Marchette, N. J.; Sidwell, R. W.; Nicholes, P. S.; and Bushman, J. B. 1962a. Studies on infectious diseases in wild animals in Utah. III. Experimental Q fever in wild vertebrates. Zoonoses Research 1: 321–339.

Marchette, N. J.; Lundgren, D. L.; Nicholes, P. S.; Bushman, J. B.; and Vest, D. 1962b. Studies on infectious diseases in wild animals in Utah. II. Susceptibility of wild mammals to experimental plague. Zoonoses Research 1: 225–229.

Marchette, N. J.; Bushman, J. B.; Parker, D. D.; and Johnson, E. E. 1962c. Studies on infectious diseases in wild animals in Utah. IV. A wild rodent (*Peromyscus* spp.) plague focus in Utah. Zoonoses Research 1: 341–361.

McAllister, C. T.; Durden, L. A.; Hnida, J. A. and Robison, H. W. 2020. Ecto- and endoparasites of the Texas deermouse, *Peromyscus attwateri*, and eastern woodrat, *Neotoma floridana* (Rodentia: Cricetidae), from Polk County, Arkansas. Journal of the Arkansas Academy of Science 74: 53–58. https://doi.org/10.54119/jaas.2020.7414

McDaniel, B. 1979. Host records of ectoparasites from small mammals of South Dakota. The Southwestern Naturalist 24: 689–691. https://doi.org/10.2307/3670531

McKenna, M. C.; and Bell, S. K. 1997. Classification of mammals above the species level. Columbia University Press, New York. 631 pp.

McLean, R. G.; Shriner, R. B.; Pokorny, K. S.; and Bowen, G. S. 1989. The ecology of Colorado tick fever in Rocky Mountain National Park in 1974. The American Journal of Tropical Medicine and Hygiene 40: 86–93. https://doi.org/10.4269/ajtmh.1989.40.86

McKeever, S. 1963. The occurrence of *Moniliformis clarki* (Ward) (Acanthocephala: Moniliformidae) in California mammals. The Journal of Parasitology 49: 1032–1033.

Meagher S. 1998. Physiological responses of deer mice (*Peromyscus maniculatus*) to infection with *Capillaria hepatica* (Nematoda). The Journal of Parasitology 84: 1112–1118.

Meagher, S. 1999. Genetic diversity and *Capillaria hepatica* (Nematoda) prevalence in Michigan deer mouse populations. Evolution 53: 1318–1324. https://doi.org/10.1111/j.1558-5646.1999.tb04547.x

Metzgar, L. H. 1971. Behavioral population regulation in the wood-mouse, *Peromyscus leucopus*. The American Midland Naturalist 86: 434–448.

Metzgar, L. H. 1973a. Comparison of trap and track-revealed home ranges in *Peromyscus*. The Journal of Mammalogy 54: 513–515.

Metzgar, L. H. 1973b. Exploratory and feeding home ranges in *Peromyscus*. The Journal of Mammalogy 54: 760–763.

Metzgar, L. H. 1979. Dispersion patterns in *Peromyscus* population. The Journal of Mammalogy 60: 129–145.

Miller, J. N. 1936. A study of *Brachylaima* (*Postharmostomum*) *sexconvolutum*, n. sp., a trematode parasite of the deer mouse. Abstract. PhD diss., Ohio State University.

Mohr, C. O.; Beck, D. E.; and Brinton, E. P. 1964. Observations on host-parasite relationships and seasonal history of ticks in San Mateo County, California. The Great Basin Naturalist 24: 1–6.

Monroe, M. C.; Morzunov, S. P.; Johnson, A. M.; Bowen, M. D.; Artsob, H.; Yates, T. L.; Peters, C. J.; Rollin, P. E.; Ksiazek, T. G.; and Nichol, S. T. 1999. Genetic diversity and distribution of *Peromyscus*-borne hantaviruses in North America. Emerging Infectious Diseases 5: 75–86. https://doi.org/10.3201/eid0501.990109

Morlan, H. B. 1954. Notes on the genus *Megarthroglossus* (Siphonaptera: Hystrichopsyllidae) in Santa Fe County, New Mexico. The Journal of Parasitology 40: 446–447.

Morlan, H. B. 1955. Mammal fleas of Santa Fe County, New Mexico. Texas Reports on Biology and Medicine 13: 93–123.

Morlan, H. B.; and Hoff, C. C. 1957. Notes on some Anoplura from New Mexico and Mexico. The Journal of Parasitology 43: 347–351.

Morshed, M. G.; Lee, M. K.; Man, S.; Fernando, K.; Wong, Q.; Hojgaard, A.; Tang, P.; Mak, S.; Henry, B.; and Patrick, D. M. 2015. Surveillance for *Borrelia burgdorferi* in *Ixodes* ticks and small rodents in British Columbia. Vector-Borne and Zoonotic Diseases 15: 701–705. https://doi.org/10.1089/vbz.2015.1854

Myton, B. 1974. Utilization of space by *Peromyscus leucopus* and other small mammals. Ecology, 55: 277–290.

Nadeau, J. H.; Lombardi, R. T.; and Tamarin, R. H. 1981. Population structure and dispersal of *Peromyscus leucopus* on Muskeget Island. Canadian Journal of Zoology 59: 793–799.

Neidert, C. M.; and Macy, R. W. 1968. *Concinnum peromysci* sp. n. (Trematoda: Dicrocoeliidae) from *Peromyscus maniculatus* in Oregon. The American Midland Naturalist 79: 525–528. https://doi.org/10.2307/2423201

Nelson, B. C.; and Smith, C. R. 1976. Ecological effects of a plague epizootic on the activities of rodents inhabiting caves at Lava Beds National Monument, California. Journal of Medical Entomology 13: 51–61. https://doi.org/10.1093/jmedent/13.1.51

Nerurkar, V. R.; Song, J. W.; Song, K. J.; Nagle, J. W.; Hjelle, B.; Jenison, S.; and Yanagihara, R. 1994. Genetic evidence for a hantavirus enzootic in deer mice (*Peromyscus maniculatus*) captured a decade before the recognition of Hantavirus Pulmonary Syndrome. Virology 204: 563–568. https://doi.org/10.1006/viro.1994.1570

Nichol, S. T.; Spiropoulou, C. F.; Morzunov, S.; Rollin, P. E.; Ksiazek, T. G.; Feldmann, H.; Sanchez, A.; Childs, J.; Zaki, S.; and Peters, C. J. 1993. Genetic identification of a hantavirus associated with an outbreak of acute respiratory illness. Science 262: 914–917. https://doi.org/10.1126/science.8235615

Nicholson, A. J. 1941. The homes and social habits of the wood mouse (*Peromyscus leucopus noveboracensis*) in southern Michigan. The American Midland Naturalist 25: 196–223.

Nieto, N. C.; Dabritz, H.; Foley, P.; Drazenovich, N.; Calder, L.; Adjemian, J.; Conrad, P. A.; and Foley, J. E. 2007. Ectoparasite diversity and exposure to vector-borne disease agents in wild rodents in central coastal California. Journal of Medical Entomology 44: 328–335. https://doi.org/10.1093/jmedent/44.2.328

Noble, G. K. 1939. The role of dominance in the social life of birds. Auk 56: 263–273.

O'Farrell, T. P. 1975. Small mammals, their parasites and pathologic lesions on the arid lands ecology reserve, Benton County, Washington. The American Midland Naturalist 93: 377–387.

Ogden C. G. 1971. Observations in the systematics of nematodes belonging to the genus *Syphacia* Seurat, 1916. Bulletin of the British Museum (Natural History), Zoology 20: 253–280.

Ohbayashi, M.; Rausch, R. L.; and Fay, F. H. 1971. On the ecology and distribution of *Echinococcus* spp. (Cestoda: Taeniidae), and characteristics of their development of larval *E. multilocularis* Leuckart, 1863, in the intermediate host. The Japanese Journal of Veterinary Research 19 (Supplement 3): 1–63.

Oliver, J.; Means, R. G.; Kogut, S.; Prusinski, M.; Howard, J. J.; Layne, L. J.; Chu, F. K.; Reddy, A.; Lee, L.; and White, D. J. 2006. Prevalence of *Borrelia burgdorferi* in small mammals in New York State. Journal of Medical Entomology 43: 924–935. https://doi.org/10.1093/jmedent/43.5.924

Oswald, V. H. 1958a. Studies on *Rictularia coloradensis* Hall, 1916 (Nematoda: Thelaziidae). I. Larval development in the intermediate host. Transactions of the American Microscopical Society 77: 229–240.

Oswald, V. H. 1958b. Studies on *Rictularia coloradensis* Hall, 1916 (Nematoda: Thelaziidae). II. Development in the definitive host. Transactions of the American Microscopical Society 77: 413–422.

Packchanian, A. 1934. Experimental *Trypanosoma brucei* infection and immunity in various species of *Peromyscus* (American deer mice). American Journal of Hygiene 20: 135–147.

Padgett, K. A.; and Boyce, W. M. 2004. Life history studies on two molecular stains of *Mesocestoides* (Cestoda: Mesocestoides): identification of sylvatic hosts and infectivity of immature life stages. The Journal of Parasitology 90: 108–113.

Page, K. L. 2013. Parasites and the conservation of small populations: The case of *Baylisascaris procyonis*. International Journal for Parasitology: Parasites and Wildlife 2: 203–210. https://doi.org/10.1016/j.ijppaw.2013.05.003

Page, L. K.; Swihart, R. K.; and Kazacos, K. R. 2001. Foraging among feces: food availability affects parasitism of *Peromyscus leucopus* by *Baylisascaris procyonis*. Journal of Mammalogy 82: 993–1002.

Padovan, D. 2006. Infectious Diseases of Wild Rodents. Corvus Publishing Company, Anacortes, Washington.

Peterson, O. 1926. The fossils of the Frankstown Cave, Blair County, Pennsylvania. Annals of the Carnegie Museum 16: 249–315.

Phan, T. G.; Kapusinszky, B.; Wang, C.; Rose, R. K.; Lipton, H. L.; and Delwart, E. L. 2011. The fecal viral flora of wild rodents. PLoS Pathogens 7. https://doi.org/10.1371/journal.ppat.1002218

Poinar, G. O.; and Nelson, B. C. 1973. *Psyllotylenchus viviparus*, n. gen., n. sp. (Nematodea: Tylenchida: Allantonematidae) parasitizing fleas (Siphonaptera) in California. Journal of Medical Entomology 10: 349–354. https://doi.org/10.1093/jmedent/10.4.349

Poirier, S. R. 1994. *Trichinella nativa* and *Trichinella pseudospiralis* in the deer mouse *Peromyscus maniculatus*: biological characterization of the infections and parasite-associated behavioral pathology of the host. PhD diss., McGill University, Montreal. https://escholarship.mcgill.ca/concern/theses/ht24wm19v

Poirier, S. R., Rau, M. E.; and Chadee, K. 1993. Enteral and parenteral phases of *Trichinella nativa* and *Trichinella pseudospiralis* in the deer mouse, *Peromyscus maniculatus*. The Journal of Parasitology 79: 733–743.

Poland, J.; Quan, T. J. and Barnes, A. M. 1994. Plague. Pp. 93–112 in: Handbook of Zoonoses. 2nd ed. Beran, G. W. (ed.) CRC Press, Boca Raton, Florida. https://doi.org/10.1201/9781003006107

Pollitzer, R.; and Meyer, K. F. 1961. The ecology of plague. Pp. 433–501 in: Studies in Disease Ecology, May, J. M. (ed.) Hafner Publishing Company, New York.

Poorbaugh, J. H.; and Gier, H. T. 1961. Fleas (Siphonaptera) of small mammals in Kansas. Journal of the Kansas Entomological Society 34: 198–204.

Porter, D. A. 1935. *Nippostrongylus muris* in the deer mouse, *Peromyscus maniculatus*. The Journal of Parasitology 21: 314.

Price, H. F. 1931. Life history of *Schistosomatium douthitti* (Cort). American Journal of Hygiene 13: 685–727.

Prince, F. M. 1944. Descriptions of three new species of *Thrassis* Jordan and the females of *T. bacchi* (Roths.) and *T. pansus* (Jordan). Pan Pacific Entomologist 20: 13–19.

Pulido-Flores, G.; Moreno-Flores, S.; and S. Monks, S. 2005. Helminths of rodents (Rodentia: Muridae) from Metztitlan, San Cristobal, and Rancho Santa Elena, Hidalgo, Mexico. Comparative Parasitology 72: 186–192.

Quan, S. F.; and Kartman, L. 1962. Ecological studies of wild rodent plague in the San Francisco Bay area of California. VIII. Susceptibility of wild rodents to experimental plague infection. Zoonoses Research 1: 121–144.

Quan, T. J.; Tsuchiya, K. R.; and Carter, L. G. 1979. Isolation of pathogens other than *Yersinia pestis* during plague investigations. Journal of Wildlife Diseases 15: 505–510. https://doi.org/10.7589/0090-3558-15.4.505

Quentin, J. C.; Karimi, Y.; and Rodriguez de Almeida, C. 1968. *Protospirura numidica criceticola* n. subsp., parasite de rongeurs Cricetidae du Brésil: cycle évolutif. Annales de Parasitologie Humaine et Comparée 43: 583–596. https://doi.org/10.1051/parasite/1968435583

Quentin, J. C.; and Kinsella, J. M. 1972. Etude de trois espèces d'oxyures *Syphacia* parasites de rongeurs cricétidés nord-américains. Annales de Parasitologie 47: 717–733.

Rand, P. W.; Lacombe, E. H.; Smith, R. P.; Rich, S. M.; Kilpatrick, W. C.; Dragoni, C. A.; and Caporale, D. 1993. Competence of *Peromyscus maniculatus* (Rodentia: Cricetidae) as a reservoir host for *Borrelia burgdorferi* (Spirochaetares: Spirochaetaceae) in the wild. Journal of Medical Entomology 30: 614–618. https://doi.org/10.1093/jmedent/30.3.614

Rankin, J. S. 1945. Ecology of helminth parasites of small mammals collected from Northrup Canyon, Upper Grand Coulee, Washington. The Murrelet 26: 11–14.

Rapp Jr., W. F. 1962. Distribution notes on parasitic mites. Acarologia 4: 31–33.

Rapp Jr., W. F.; and Gates, D. B. 1957. A distributional checklist of the fleas of Nebraska. Journal of the Kansas Entomological Society 30: 50–53.

Rausch, R. 1949. Observations of the life cycle and larval development of *Paruterina candelabraria* (Goeze, 1782) (Cestoda: Dilepididae). The American Midland Naturalist 42: 713–721.

Rausch, R. 1961. Notes on the occurrence of *Capillaria hepatica* (Bancroft, 1893). Proceedings of the Helminthological Society of Washington 28: 17–18.

Rausch, R. L. 1967. On the ecology and distribution of *Echinococcus* spp. (Cestoda: Taeniidae) and characteristics of their development in the intermediate host. Annales de Parasitologie Humaine et Comparée 42: 19–62.

Rausch, R. L; and Richards, S. H. 1971. Observations on parasite-host relationships of *Echinococcus multilocularis* Leuckart, 1863 in North Dakota. Canadian Journal of Zoology 49: 1317–1330.

Rausch, R.; and Schiller, E. L. 1954. Studies on the helminth fauna of Alaska. XXIV. *Echinococcus sibiricensis* n. sp. from St. Lawrence Island. The Journal of Parasitology 40: 659–662.

Read, C. P. 1949a. Studies on North American helminths of the genus *Capillaria* Zeder, 1800 (Nematoda) II. Additional capillarids from mammals with keys to the North American mammalian species. The Journal of Parasitology 35: 231–239.

Read, C. P. 1949b. Studies on North American helminth of the genus *Capillaria* Zeder, 1800 (Nematoda) I. *Capillaria* from mammals. The Journal of Parasitology 35: 223–230.

Redetzke, K. A.; and Canaris, A. G. 1977. *Brachylaime microti*: A mechanistic simulation model of the parasite, its intermediate snail host, *Oreohelix strigosa*, and its definitve rodent hosts, *Peromyscus maniculatus* and *Microtus montanus*. Experimental Parasitology 41: 229–271.

Redetzke, K. A.; and McCann, M. J. 1980. Isolation of *Leptospira* from desert rodents of west Texas. Journal of Wildlife Diseases 16: 333–337.

Reduker, D. W.; Hertel, L.; and Duszynski, D. W. 1985. *Eimeria* species (Apicomplexa: Eimeriidae) infecting *Peromyscus* rodents in the southwestern United States and northern Mexico with description of a new species. The Journal of Parasitology 71: 604–613. https://doi.org/10.2307/3281432

Reduker, D. W.; Duszynski, D. W.; and Yates, T. L. 1987. Evolutionary relationships among *Eimeria* spp. (Apicomplexa) infecting cricetid rodents. Canadian Journal of Zoology 65: 722–735.

Redman, J. P.; and Sealander, J. A. 1958. Home ranges of deer-mice in southern Arkansas. The Journal of Mammalogy 30: 390–395.

Reed, J. T. 1973. *Comatacarus*, formerly subgenus of *Leeuwenhoekia*, restored to generic status, with description of a new species (Acarina: Trombiculidae). Journal of Medical Entomology 10: 315–317. https://doi.org/10.1093/jmedent/10.3.315

Reeves, W. C.; Scrivani, R. P.; Puge, W. E.; and Rowe, W. P. 1967. Recovery of an adenovirus from a feral rodent *Peromyscus maniculatus*. Proceedings of the Society for Experimental Biology and Medicine 124: 1173–1175. https://doi.org/10.3181/00379727-124-31955

Reeves, W. K.; Durden, L. A.; Ritzi, C. M.; Beckham, K. R.; Super, P. E. and OConnor, B. M. 2007. Ectoparasites and other ectosymbiotic arthropods of vertebrates in the Great Smoky Mountains National Park, USA. Zootaxa 1392: 31–68.

Richard, W. H. 1960. The distribution of small mammals in relation to the climax vegetation mosaic in eastern Washington and northern Idaho. Ecology 41: 99–106.

Riser, N. W. 1956. The hooks of taenioid cestodes from North American felids. The American Midland Naturalist 56: 133–137.

Robert, A. 1962. Siphonapteres récoltés sur les petites rongeurs du Parc du Mont Tremblant, et leure relations avec leur hôtes. Annals of the Entomological Society of Québec 7: 3–18.

Robinson, E. J. 1949. The life history of *Postharmostomum helicis* (Leidy, 1847) n. comb. (Trematoda: Brachylaemidae). The Journal of Parasitology 35: 513–533.

Rodenberg, G. W.; and Pence, D. B. 1978. Circulation of helminth species in a rodent population from the high plains of Texas. Occasional Papers of the Museum Texas Tech University 56: 1–10.

Rodríguez-Rojas, J. J.; Rodríguez-Moreno, Á.; Sánchez-Casas, R. M.; and Hernández-Escareño, J. J. 2020. Molecular detection of *Leptospira interrogans* and *Borrelia burgdorferi* in wild rodents from Mexico. Vector-Borne and Zoonotic Diseases 20: 860–863. https://doi.org/10.1089/vbz.2019.2600

Root, J. J.; Calisher, C. H.; and Beaty, B. J. 1999. Relationships of deer mouse movement, vegetative structure, and prevalence of infection with Sin Nombre virus. Journal of Wildlife Diseases 35: 311–318. https://doi.org/10.7589/0090-3558-35.2.311

Rothschild, N. C. 1906. Three new Canadian fleas. The Canadian Entomologist 38: 321–325.

Rupes, V.; and Whitaker Jr., J. O. 1968. Mites of the subfamily Labidophorinae (Acaridae, Acarina) in North America. Acarologia 10: 493–499.

Rutledge, L. C.; Moussa, M. A.; Zeller, B. L.; and Lawson, M. A. 1979. Field studies of reservoirs and vectors of sylvatic plague at Fort Hunter Liggett, California. Journal of Medical Entomology 15: 452–458. https://doi.org/10.1093/jmedent/15.5-6.452

Safronetz, D.; Drebot, M. A.; Artsob, H.; Cote, T.; Makowski, K.; and Lindsay, L. R. 2008. Sin Nombre virus shedding patterns in naturally infected deer mice (*Peromyscus maniculatus*) in relation to duration of infection. Vector-Borne and Zoonotic Diseases 8: 97–100. https://doi.org/10.1089/vbz.2007.0113

Sapp, S. G. H.; Rascoe, L. N.; Wilkins, P. P.; Handali, S.; Gray, E. B.; Eberhard, M.; Woodhall, D. M.; Montgomery, S. P.; Bailey, K. L.; Lankau, E. W.; and Yabsley, M. J. 2016a. *Baylisascaris procyonis* roundworm seroprevalence among wildlife rehabilitators, United States and Canada, 2012–2015. Emerging Infectious Diseases 22: 2128–2131. https://doi.org/10.3201/eid2212.160467

Sapp, S. G.; Weinstein, S. B.; McMahan, C. S.; and Yabsley, M. J. 2016b. Variable infection dynamics in four *Peromyscus* species following experimental inoculation with *Baylisascaris procyonis*. The Journal of Parasitology 102: 538–544.

Sapp, S. G.; Handali, S.; Weinstein, S. B.; and Yabsley, M. J. 2018. Detection and evaluation of antibody response to a *Baylisascaris*-specific antigen in rodent hosts with the use of Western blotting and ELISA. The Journal of Parasitology 104: 651–659.

Scanlon, J. E. 1960. The Anoplura and Mallophaga of the mammals of New York. Wildlife Diseases No 5. (Printed on 3 microcards.)

Schad, G. A. 1954. Helminth parasites of mice in northeastern Quebec and the Coast of Labrador. Canadian Journal of Zoology 32: 215–224.

Schad, G. A. 1956. Helminths recovered from deer mice, *Peromyscus maniculatus*, trapped in the Morgan Arboretum, MacDonald College, Provence of Quebec. Canadian Journal of Zoology 34: 208.

Scharf, W. C.; and Stewart, K. R. 1980. New records of Siphonaptera from northern Michigan. Great Lakes Entomologist 13: 165–167.

Scherer, W. F.; Dickerman, R. W.; La Fiandra, R. P.; Wong Chia, C.; and Terrian, J. 1971. Ecologic studies of Venezuelan encephalitis virus in southeastern México. IV. Infections of

wild mammals. The American Journal of Tropical Medicine and Hygiene 20: 980–988. https://doi.org/10.4269/ajtmh.1971.20.980

Schmaljohn, A. L.; Li, D.; Negley, D. L.; Bressler, D. S.; Turell, M. J.; Korch, G. W.; Ascher, M. S.; and Schmaljohn, C. S. 1995. Isolation and initial characterization of a newfound hantavirus from California. Virology 206: 963–972. https://doi.org/10.1006/viro.1995.1019

Schneider, B. S.; Zeidner, N. S.; Burkot, T. R.; Maupin, G. O.; and Piesman, J. 2000. *Borrelia* isolates in northern Colorado identified as *Borrelia bissettii*. Journal of Clinical Microbiology 38: 3103–3105.

Scholten, T. H.; Ronald, K.; and McLean, D. M. 1962. Parasite fauna of the Manitoulin Island Region. I. Arthropoda parasitica. Canadian Journal of Zoology 40: 605–606.

Schwan, T. C.; and Dobkin, D. S. 1981. An unusual example of teratogenesis in the flea *Thrassis fotus* from Colorado (Siphonaptera: Ceratophyllidae). Proceedings of the Entomological Society of Washington 83: 93–98.

Schwanz, L. E. 2006. Schistosome infection in deer mice (*Peromyscus maniculatus*): impacts on host physiology, behavior and energetics. Journal of Experimental Biology 209: 5029–5037.

Seaman, R. N.; and Nash, D. J. 1976. Variation in the incidence of botfly larvae (*Cuterebra*) in two sympatric species of *Peromyscus* in northern Colorado. Great Basin Naturalist 36: 481–482.

Semtner, P. J.; and Hair, J. A. 1973. Distribution, seasonal abundance, and hosts of the Gulf Coast tick in Oklahoma. Annals of the Entomological Society of America 66: 1264–1268. https://doi.org/10.1093/aesa/66.6.1264

Senger, C. M.; and Macy, R. W. 1952. Helminths of northwest mammals. Part III. The description of *Euryhelmis pacificus* n. sp. and notes on its life cycle. The Journal of Parasitology 38: 481–486.

Senger, C. M. 1966. Notes of Fleas (Siphonaptera) from Montana. Journal of the Kansas Entomological Society 39: 105–109.

Seymour, C.; and Yuill, T. 1981. Arboviruses [Arthropod-borne virus]. Pp. 54–86 in: Infectious Diseases of Wild Mammals, Karastad, L. H.; Davis, J. W.; and Trainer, D. O. (eds.). Iowa State University Press, Ames.

Shaftesbury, A. D. 1934. The Siphonaptera (fleas) of North Carolina with special reference to sex ratios. Journal of the Elisha Mitchell Scientific Society 49: 247–263.

Sheppe, W. 1966. Determinants of home range in the deer mouse, *Peromyscus leucopus*. Proceedings of the California Academy of Sciences 34: 377–418.

Short, R. B. 1951. Hermaphroditic female *Schistosomatium douthitti* (Trematoda: Schistosomatidae). The Journal of Parasitology 37: 547–555.

Short, R. B. 1952. Sex studies on *Schistosomatium douthitti* (Cort, 1914) Price, 1931 (Trematoda: Schistosomatidae). The American Midland Naturalist 47: 1–54.

Sidwell, R. W.; Lundgren, D. L.; Bushman, J. B. and Thorpe, B. D. 1964. The occurrence of a possible epizootic of Q fever in fauna of the Great Salt Lake desert of Utah. The American Journal of Tropical Medicine and Hygiene 13: 754–762.

Sigler, L. 2005. Adiaspiromycosis and other infections caused by *Emmonsia* species. Pp. 810–824 in: Topley and Wilson's microbiology and microbial infections, 10th ed., Cox, F. E. G.; Wakelin, D.; Gillespie, S.; and Despommier, D. D. (eds.). ASM Press, Washington, DC.

Simpson, D. 1968. Arboviruses and free living wild animals. Pp. 13–28 in: Symposia of the Zoological Society of London, Volume 24, McDiarmid, A. (ed.). Academic Press, London, UK.

Skinker, M. S. 1935. Two new species of tapeworms from carnivores and a redescription of *Taenia laticolis* Rudolphi, 1819. Proceedings of the United States National Museum 83: 211–220.

Slaughter, B. H.; and Ritchie, R. 1963. Pleistocene mammals of the Clear Creek local fauna, Denton County, Texas. Southern Methodist Journal of the Graduate Research Center 31: 117–131. Cited by Hibbard (1968).

Smith, C. F. 1953. Studies on the helminth fauna of small mammals in the areas of the High Plains, central and southern Rockies. PhD diss., University Nebraska, Lincoln.

Smith, C. F. 1954. Four new species of cestodes of rodents from the High Plains, central and southern Rockies and notes on *Catenotaenia dendritica*. The Journal of Parasitology 40: 245–254.

Smith, D. H. 1975. An ecological analysis of a host parasite association: *Cuterebra approximata* (Diptera: Cuterebridae) in *Peromyscus maniculatus* (Rodentia: Cricetidae). PhD diss., University of Montana, Missoula.

Smith, D. H. 1977. The natural history and development of *Cuterebra approximata* (Diptera: Cuterebridae) in its natural host, *Peromyscus maniculatus* (Rodentia: Cricetidae), in western Montana. Journal of Medical Entomology 14: 137–145. https://doi.org/10.1093/jmedent/14.2.137

Smith, K. F.; and Carpenter, S. M. 2006. Spread of introduced black rat (*Rattus rattus*) parasites to endemic deer mice (*Peromyscus maniculatus*) on the California Channel Islands. Diversity and Distributions 12: 742–748.

Solomon, G. B.; and Warner, G. S. 1969. *Trichinella spiralis* in mammals at Mountain Lake, Virginia. The Journal of Parasitology 55: 730–732.

Solomon, G. B.; and Handley Jr., C. O. 1971. *Capillaria hepatica* (Bancroft, 1893) in Appalachian mammals. The Journal of Parasitology 57: 1142–1144. https://doi.org/10.2307/3277883

Sonenshine, D. E.; Yunker, C. E.; Clifford, C. M.; Clark, G. M. and Rudbach, J. A. 1976. Contributions to the ecology of Colorado Tick Fever Virus 2: Population dynamics and host utilization of immature stages of the Rocky Mountain wood tick, *Dermacentor andersoni*. Journal of Medical Entomology 12: 651–656. https://doi.org/10.1093/jmedent/12.6.651

Song, J. W.; Baek, L. J.; Nagle, J. W.; Schlitter, D.; and Yanagihara, R. 1996. Genetic and phylogenetic analyses of hantaviral sequences amplified from archival tissues of deer mice (*Peromyscus maniculatus nubiterrae*) captured in the eastern United States. Archives of Virology 141: 959–967. https://doi.org/10.1007/BF01718170

Spencer, C. J. 1940. Ectoparasites of birds and mammals in British Columbia. VI. Perlimiary list of parasitic mites. Proceedings of the Entomological Society of British Columbia 37: 14–18.

Spencer, G. J. 1966. Anoplura from British Columbia and some adjacent areas. Journal of the Entomological Society of British Columbia 63: 23–30.

Stallard, H. E.; and Arai, H. P. 1978. The growth and development of *Hymenolepis peromysci* Tinkle, 1972 (Cestoda: Cyclophyllidea). Canadian Journal of Zoology 56: 90–93.

Stanford, J. S. 1931. A preliminary list of Utah Siphonaptera. Proceedings Utah Academy Sciences 8: 153–154.

Stapp, P.; Salkeld, D. J.; Eisen, R. J.; Pappert, R.; Young, J.; Carter, L. G.; Gage, K. L.; Tripp, D. W.; and Antolin, M. F. 2008. Exposure of small rodents to plague during epizootics in black-tailed prairie dogs. Journal of Wildlife Diseases 44: 724–730. https://doi.org/10.7589/0090-3558-44.3.724

Stark, H. E. 1958. The Siphonaptera of Utah. U.S. Department of Health, Education and Welfare, Communicable Disease Center. Atlanta, Georgia. 239 pp.

Stark, H. E.; and Kinney, A. R. 1969. Abundance of rodents and fleas as related to plague in Lava Beds National Monument, California. Journal of Medical Entomology 6: 287–294.

Stark, H. E.; Campos, E. G.; and Elbel, R. E. 1976. Description of the third-instar larva of *Monopsyllus wagneri* (Baker) (Siphonaptera: Ceratophyllidae). Journal of Medical Entomology 13: 107–111. https://doi.org/10.1093/jmedent/13.1.107

Steere, A. C. 1989. Lyme disease. New England Journal of Medicine 321: 586–596. https://doi.org/10.1056/NEJM198908313210906

Stewart. M. A. 1933. Revision of the list of Siphonaptera from New York State. Journal of the New York Entomological Society 41: 253–262.

Stickel, L. F. 1960. *Peromyscus* ranges at high and low densities. The Journal of Mammalogy 41: 433-441.

Stock, A. D. 1961. Endoparasites of mammals found in the Navajo Reservoir basin. Pp. 91–94 in: University of Utah Anthropological Papers, No. 55; Upper Colorado Series, No. 5. University of Utah Press, Salt Lake City.

Stock, A. D. 1962. Endoparasites of mammals from the Curecanti Reservoir Basins, western Colorado. Pp. 162–166 in: University of Utah Anthropplogical Papers, No. 59; Upper Colorado Series, No. 8. University of Utah Press, Salt Lake City.

Stoenner, H. G.; Holdenried, R.; Lackman, D.; and Osborn Jr., J. S. 1959. The occurrence of *Coxiella burnetii*, *Brucella*, and other pathogens among fauna of the Great Salt Lake Desert in Utah. The American Journal of Tropical Medicine and Hygiene 8: 590–596. https://doi.org/10.4269/ajtmh.1959.8.590

Storm, J. J.; and Ritzi, C. M. 2008. Ectoparasites of small mammals in western Iowa. Northeastern Naturalist 15: 283–292. https://doi.org/10.1656/1092-6194(2008)15[283:EOSMIW]2.0.CO;2

Strandtman, R. W.; and Allred, D. M. 1956. Mites of the genus *Brevisguma* Keegan, 1949 (Acarina: Haemogamasidae). Journal of the Kansas Entomological Society 29: 113–132.

Strandtman, R. W.; and Morlan, H. B. 1953. A new species of *Hirstionyssus* and a key to the known species of the World. Texas Reports on Biology and Medicine 11: 627–637.

Sudia, W. D.; and Newhouse, V. F. 1975. Epidemic Venezuelan equine encephalitis in North America: a summary of virus-vector-host relationships. American Journal of Epidemiology 101: 1–13.

Swanekamp, L. J. 2005. The role of rodents as a potential reservoir for *Pasteurella multocida* on the National Elk Refuge, Wyoming. Master's thesis, Montana State University, Bozeman. 61 pp. https://scholarworks.montana.edu/xmlui/handle/1/2381

Taitt, M. J. 1981. The effect of extra food on small rodent populations. I. Deermice (*Peromyscus maniculatus*). The Journal of Animal Ecology 50: 111–124.

Tamsitt, J. R. 1957. *Peromyscus* from the late Pleistocene of Texas. Texas Journal of Science 9: 355–363. Cited by Hibbard (1968).

Tanigoshi, L. K.; and Loomis, R. B. 1974. The genus *Hyponeocula* (Acarina, Trombiculidae) of western North America. Melanderia 17: 1–27.

Taylor, R. M. 1967. Catalogue of arthropod-borne viruses of the world. US Department of Health, Education, and Welfare, Public Health Service.

Terman, C. R. 1961. Some dynamics of spatial distribution within semi-natural populations of prairie deermice. Ecology 42: 288–302.

Test, F. H.; and Test, A. R. 1943. Incidence of dipteran parasitosis in populations of small mammals. The Journal of Mammalogy 24: 506–508.

Theis, J. H.; and Schwab, R. G. 1992. Seasonal prevalence of *Taenia taeniaeformis*: relationship to age, sex, reproduction and abundance of an intermediate host (*Peromyscus maniculatus*). Journal of Wildlife Diseases 28: 42–50.

Thompson, M.; Mykytczuk, N.; Gooderham, K.; and Schulte-Hostedde, A. 2012. Prevalence of the bacterium *Coxiella burnetii* in wild rodents from a Canadian natural environment park. Zoonoses and Public Health 59: 553–560. https://doi.org/10.1111/j.1863-2378.2012.01493.x

Tiner, J. D. 1948a. *Syphacia eutamii* n. sp. from the least chipmunk, *Eutamias minimus*, with a key to the genus (Nematoda: Oxyuridae). The Journal of Parasitology 34: 87–92.

Tiner, J. D. 1948b. Observations of the *Rictularia* (Nematoda: Thelaziidae) of North America. Transactions of the American Microscopical Society 67: 192–200.

Tiner, J. D. 1953. Fatalities in rodents caused by larval *Ascaris* in the central nervous system. The Journal of Mammalogy 34: 15.

Tiner, J. D.; and Rausch, R. 1950. Two new *Syphacia* (Nematoda: Oxyuridae) and observations on the inner circle circumoral papillae in North American species of the genus. Chicago Academy of Sciences Natural History Miscellanea 57: 1048–1051.

Tinkle, D. P. 1972. Description and natural intermediate hosts of *Hymenolepis peromysci* n. sp., a new cestode from deer mice (*Peromyscus*). Transactions of the American Microscopical Society 91: 66–69.

Tipton, V. J.; and Mendez, E. 1968. New species of fleas (Siphonaptera) from Cerro Potosi, Mexico, with notes on ecology and host parasite relationships. Pacific Insects 10: 177–214.

Traub, R.; and Nadchatram, N. 1966. A revision of the genus *Chatia* Brennan, with synonymic notes and descriptions of two new species from Pakistan (Acarina: Trombiculidae). Journal of Medical Entomology 2: 373–383.

Travassos, L.; and Darriba, A. R. 1929. Notas sobre Heligmosominae. Sciencia Medica. 7: 432–438.

Twigg, G. I. 1978. The role of rodents in plague dissemination: a worldwide review. Mammal Review 8: 77–110.

Tyzzer, E. E. 1942. A comparative study of Grahamellae, Haemobartonellae and Eperythrozoa in small mammals. Proceedings of the American Philosophical Society 85: 359–398.

Ubelaker, J. E.; Abdullah, N.; Mouhaffel, A.; Ananadampillair, R.; Emigh, C.; and Gardner, S. L. 2014. Natural infections of tetrathyridia of *Mesocestoides* species in deer mice, *Peromyscus maniculatus*, from New Mexico. The Southwestern Naturalist 59: 404–406. https://doi.org/10.1894/JC-80.1

Ubico, S. R.; McLean, R. G.; and Cooksey, L. M. 1996. Susceptibility of selected rodent species from Colorado to *Borrelia burgdorferi*. Journal of Wildlife Diseases 32: 293–299. https://doi.org/10.7589/0090-3558-32.2.293

Ulmer, M. J. 1951. *Postharmostomatum helicis* (Leidy, 1847) Robinson, 1949 (Trematoda). Its life history and a revision of the family Brachylaeminae. Part 1. Transactions of the American Microscopical Society 70: 189–238.

Ulrich, M. G.; and Vaughn, C. M. 1963. Some intestinal and external parasites of the deer mouse, *Peromyscus maniculatus*. Proceedings of the South Dakota Academy of Sciences 42: 140–143.

Van Cleave, H. J. 1953. Acanthocephala of North American Mammals. Illinois Biological Monographs, Vol. 23, Nos. 1–2. The University of Illinois Press, Urbana. 179 pp.

Van Horne, B. 1982. Niches of adult and juvenile deer mice, *Peromyscus manculatus*, in seral stages of coniferous forest. Ecology 63: 922–1003.

Vanderwolf, K. J.; Malloch, D.; and McAlpine, D. F. 2018. Psychrotolerant microfungi associated with deer mice (*Peromyscus maniculatus*) in a white-nose syndrome positive bat hibernaculum in eastern Canada. The Canadian Field-Naturalist 131: 238–245. https://doi.org/10.22621/cfn.v131i3.1906

Vercammen-Grandjean, P. 1965. Revision of the genera: *Eltonella* Audy, 1956 and *Microtrombicula* Ewing, 1950, with descriptions of fifty new species and transferal of subgenus *Chiroptella* to genus *Leptotrombidium* (Acarina, Trombiculidae). Acarologia 7: 34–257.

Verma, A.; Beigel, B.; Smola, C. C.; Kitts-Morgan, S.; Kish, D.; Nader, P.; Morgan, J.; Roberson, J.; Christmann, U.; Gruszynski, K.; Brandt, L.; Cho, E.; Murphy, K.; and Goss, R. 2019. Evidence of leptospiral presence in the Cumberland Gap region. PLOS Neglected Tropical Diseases 13: e0007990. https://doi.org/10.1371/journal.pntd.0007990

Verts, B. J. 1961. Observations of the fleas (Siphonaptera) of some small mammals in northwestern Illinois. American Midland Naturalist 66: 471–476.

Vest, E. D.; and Marchette, N. J. 1958. Transmission of *Pasteurella tularensis* among desert rodents through infective carcasses. Science 128: 363–364.

Vest, E. D.; Lundgren, D. L.; Parker, D. D.; Johnson, D. E.; Morse, E. L.; Bushman, J. B.; Sidwell, R. W.; and Thorpe, B. D. 1965. Results of a five-year survey for certain enzootic diseases in the fauna of western Utah. The American Journal of Tropical Medicine and Hygiene 14: 124–135. https://doi.org/10.4269/ajtmh.1965.14.124

Voge, M. 1955. A list of cestode parasites from California mammals. American Midland Naturalist 54: 413–417.

Vredevoe, L. K.; Stevens, J. R.; and Schneider, B. S. 2004. Detection and characterization of *Borrelia bissettii* in rodents from the central California coast. Journal of Medical Entomology 41: 736–745. https://doi.org/10.1603/0022-2585-41.4.736

Wagner, J. 1936. The fleas of British Columbia. The Canadian Entomologist 68: 193–207.

Wassom, D. L.; Guss, V. M.; and Grundmann, A. W. 1973. Host resistance in a natural host-parasite system. Resistance to *Hymenolepis citelli* by *Peromyscus maniculatus*. The Journal of Parasitology 59: 117–121. https://doi.org/10.2307/3278585

Wassom, D. L.; DeWitt, C. W.; and Grundmann, A. W. 1974. Immunity to *Hymenolepis citelli* by *Peromyscus maniculatus*: genetic control and ecological implications. The Journal of Parasitology 60: 47–52. https://doi.org/10.2307/3278676

Wassom, D. L.; Dick, T. A.; Arnason, N.; Strickland, D.; and Grundmann, A. W. 1986. Host genetics: a key factor in regulating the distribution of parasites in natural host populations. The Journal of Parasitology 72: 334–337. https://doi.org/10.2307/3281615

Watson, E. A.; and Hadwen, S. 1912. Trypanosomes found in Canadian mammals. Parasitology 5: 11–21.

Webb Jr., J. P., and Loomis, R. B. 1970. A new subgenus of intranasal chiggers of the genus *Microtrombicula* from North America and Korea. Journal of Medical Entomology 7: 655–663. https://doi.org/10.1093/jmedent/7.6.655

Whitaker Jr., J. O. 1966. Food of *Mus musculus*, *Peromyscus maniculatus bairdi* and *Peromyscus leucopus* in Vigo County, Indiana. Journal of Mammalogy 47: 473–486. https://doi.org/10.2307/1377688

Whitaker Jr., J. O. 1968. Parasites. Pp. 254–311 in: Biology of *Peromyscus*. King, J. (ed.). American Society of Mammalogists, Stillwater, Oklahoma.

Whitaker Jr., J. O.; and Corthum Jr., K. W. 1967. Fleas of Vigo County, Indiana. Proceedings of the Indiana Academy of Science 76: 431–440.

Whitaker Jr., J. O.; and Ferraro, M. G. 1963. Summer food of 220 short-tailed shrews from Ithaca, New York. Journal of Mammalogy 44: 419. https://doi.org/10.2307/1377216

Whitaker, J. O.; and Wilson, N. 1968. Mites of small mammals of Vigo County, Indiana. The American Midland Naturalist 80: 537–542. https://doi.org/10.2307/2423545

Whitaker, J. O.; Jones, G. S.; and Pascal, D. D. 1975. Notes on mammals of the Fires Creek Area, Nantahala Mountains, North Carolina, including their ectoparasites. Journal of the Elisha Mitchell Scientific Society 91: 13–17.

Whitaker Jr., J. O.; and Maser, C. 1985. Mites (excluding chiggers) of mammals of Oregon. The Great Basin Naturalist 45: 67–76.

Whitney, E.; Jamnback, H.; Means, R. G.; and Watthews, T. H. 1968. Arthropod-borne-virus survey in St. Lawrence County, New York: arbovirus reactivity in serum from amphibians, reptiles, birds, and mammals. The American Journal of Tropical Medicine and Hygiene 17: 645–650. https://doi.org/10.4269/ajtmh.1968.17.645

Williams, L. A.; and Hoff, C. C. 1951. Fleas from the upper Sonoran zone near Albuquerque, New Mexico. Proceedings of the United States National Museum 101: 305–313.

Wilson, N. 1957. Some ectoparasites from Indiana mammals. Journal of Mammalogy 38: 281–282. https://doi.org/10.2307/1376342

Wilson, N. 1961. The ectoparasites (Ixodides, Anoplura and Siphonaptera) of Indiana mammals. Master's thesis, Purdue University, West Lafayette, Indiana. Cited by Whitaker (1968).

Wobeser, G.; Ngeleka, M.; Appleyard, G.; Bryden, L.; and Mulvey, M. R. 2007. Tularemia in deer mice (*Peromyscus maniculatus*) during a population irruption in Saskatchewan, Canada. Journal of Wildlife Diseases 43: 23–31. https://doi.org/10.7589/0090-3558-43.1.23

Wobeser, G.; Campbell, G. D.; Dallaire, A.; and McBurney, S. 2009. Tularemia, plague, yersiniosis, and Tyzzer's disease in wild rodents and lagomorphs in Canada: a review. The Canadian Veterinary Journal 50: 1251–1256.

Wolfenbarger, K. A. 1952. Systematic and biological studies on North American chiggers of the genus *Trombicula*, subgenus *Eutrombicula* (Acarina, Trombiculidae). Annals of the Entomological Society of America 45: 645–677. https://doi.org/10.1093/aesa/45.4.645

Wolff, J. O. 1985a. Comparative population ecology of *Peromyscus leucopus* and *Peromyscus maniculatus*. Canadian Journal of Zoology 63: 1548–1555. https://doi.org/10.1139/z85-230

Wolff, J. O. 1985b. The effects of density, food, and interspecific interference on home range size in *Peromyscus leucopus* and *Peromyscus maniculatus*. Canadian Journal of Zoology 63: 2657–2662. https://doi.org/10.1139/z85-397

Wolff, J. O.; and Lundy, K. I. 1985. Intra-familial dispersion patterns in white-footed mice, *Peromyscus leucopus*. Behavioral Ecology and Sociobiology 17: 381–384. https://doi.org/10.1007/BF00293216

Wolff, J. O. 1993. Why are female small mammals territorial? Oikos 68: 364–370. https://doi.org/10.2307/3544853

Wolfgang, R. W. 1956. Helminth parasites of reptiles, birds, and mammals in Egypt. II. *Catenotaenia aegyptica* sp. nov. from myomorph rodents, with additional notes on the genus. Canadian Journal of Zoology 34: 6–20. https://doi.org/10.1139/z56-002

Wood, F. D. 1934. Natural and experimental infection of *Triatoma protracta* Ubler and mammals in California with American human trypanosomiasis. American Journal of Tropical Medicine 14: 497–511.

Wood, S. F. 1942. Observations on Chagas' disease in the United States. I. California. Bulletin of the Southern California Academy of Science 41: 61–69.

Wood, S. F. 1952. Mammal blood parasite records from southwestern United States and Mexico. The Journal of Parasitology 38: 85–86.

Wood, S. F. 1962. Blood parasites of mammals of the Californian Sierra Nevada foothills, with special reference to *Trypanosoma cruzi* Chagas and *Hepatozoon leptosoma* sp. n. Bulletin of the Southern California Academy of Sciences 61: 161–176.

Wood, S. F. 1975. New localities for mammal blood parasites from southwestern United States. The Journal of Parasitology 61: 969–970.

Woods, C. E.; and Larson, O. R. 1970. North Dakota fleas. II. Records from man and other mammals. Proceedings of the North Dakota Academy of Science 23: 31–40.

Woods, C. E.; and Larson, O. R. 1971. North Dakota fleas. III. Additional records from mammals. Proceedings of the North Dakota Academy of Science 24: 36–29.

Worley, D. E.; Meisenhelder, J. E.; Sheffield, H. G.; and Thompson, P. E. 1960. Laboratory studies of the rodent whipworm, *Trichuris muris* (Schrank, 1788). The Journal of Parasitology 46: 38.

Wrenn, W. J.; and Loomis, R. B. 1973. A new species of *Euschoengastia* (Acarina: Trombiculidae) from western North America and the status of *E. californica* (Ewing). Journal of Medical Entomology 10: 97–100. https://doi.org/10.1093/jmedent/10.1.97

Wright, K. A. 1961. Observations on the life cycle of *Capillaria hepatica* (Bancroft, 1893) with a description of the adult. Canadian Journal of Zoology 39: 167–182. https://doi.org/10.1139/z61-022

Wyman, R. L.; and Schaefer, K. 1972. The ectoparasites of *Peromyscus maniculatus* in Texas County, Oklahoma. Southwestern Naturalist 16: 435–436.

Yamaguti, S. 1958–1963. Systema Helminthum. 5 volumes. Interscience Publishers, New York.

Zarnke, R. L.; and Yuill, T. M. 1985. Modoc-like virus isolated from wild deer mice (*Peromyscus maniculatus*) in Alberta. Journal of Wildlife Diseases 21: 94–99. https://doi.org/10.7589/0090-3558-21.2.94

Zeidner, N. S.; Burkot, T. R.; Massung, R.; Nicholson, W. L.; Dolan, M. C.; Rutherford, J. S.; Biggerstaff, B. J.; and Maupin, G. O. 2000. Transmission of the agent of human granulocytic ehrlichiosis by *Ixodes spinipalpis* ticks: evidence of an enzootic cycle of dual infection with *Borrelia burgdorferi* in northern Colorado. The Journal of Infectious Diseases 182: 616–619. https://doi.org/10.1086/315715

Zenchak, J. J.; and Hall, J. E. 1971. Helminths from *Peromyscus leucopus* and *P. maniculatus* in West Virginia. The Journal of Parasitology 57: 542.

Zwolak, R.; Meagher, S.; Vaughn, J. W.; Dziemian, S.; and Crone, E. E. 2013. Reduced ectoparasite loads of deer mice in burned forest: from fleas to trees? Ecosphere 4: art132. https://doi.org/10.1890/ES13-00138.1

Peromyscus maniculatus. *National Park Service photo by John Good.*